

WZ Sge-Type Dwarf Novae

Taichi KATO^{1*}

¹ *Department of Astronomy, Kyoto University, Kyoto 606-8502*

**tkato@kusastro.kyoto-u.ac.jp*

(Received 201 0; accepted 201 0)

Abstract

We have summarized the current understanding and recently obtained findings about WZ Sge-type dwarf novae. We also reviewed the historical development of the understanding of these objects, provided the modern criteria, and reviewed the past research in relation to superhumps, early superhumps and the outburst mechanism. We regard that the presence of early superhumps (reflecting the 2:1 resonance) and long or multiple rebrightenings are the best distinguishing properties of WZ Sge-type dwarf novae. We provided the updated list of nearly 100 WZ Sge-type dwarf novae mainly based on the data obtained by the VSNET Collaboration up to Kato et al. (2015) and discussed the statistics. We could detect early superhumps with amplitude larger than 0.02 mag in 63% of the studied WZ Sge-type dwarf novae, which makes early superhumps a useful distinguishing feature for WZ Sge-type dwarf novae. Theoretical light curves of early superhumps generally appear to reproduce the existence of many low-amplitude objects, supporting the geometrical origin of early superhumps. Using the recently developed method of measuring mass ratios using developing phase of superhumps (stage A superhumps), we showed that there is a linear relation between the period variation of superhumps and the mass ratio in WZ Sge-type objects. By using this relation, we were able to draw an evolutionary picture of a large number of WZ Sge-type and identified the type of outburst to be an evolutionary sequence: type C \rightarrow D \rightarrow A \rightarrow B \rightarrow E, with some outliers for type-B objects. The duration of stage A (evolutionary phase) of superhumps is also well correlated with the estimated mass ratios. By using mass ratios from stage A superhumps and duration of stage A, we have been able to identify best candidates for period bouncers.

Key words: accretion, accretion disks — stars: novae, cataclysmic variables — stars: dwarf novae — stars: evolution — surveys

1. Introduction

“How many WZ Sge-type dwarf novae are known?” — this recent question by an acquainted astronomer was the motivation of this paper.

WZ Sge-type dwarf novae (DNe) or WZ Sge-type objects are a class of SU UMa-type dwarf novae which is a kind of cataclysmic variables (CVs) [For general information of CVs, DNe, SU UMa-type dwarf novae and superhumps, see e.g. Warner (1995)]. SU UMa-type dwarf novae are defined by the presence of superhumps, which have periods longer than the orbital period by a few percent. SU UMa-type dwarf novae are known to show short, normal outbursts and superoutbursts and superhumps are always present during superoutbursts. WZ Sge-type dwarf novae were originally proposed as a group of dwarf novae with unusually high (~ 8 mag) amplitudes and rare outbursts (approximately once a decade). The definition has been changing as our knowledge improved, and currently the definition is somewhat different from the original one.

In the 1990s, there were only a handful of WZ Sge-type dwarf novae and they were generally considered to be a rare population of dwarf novae or cataclysmic variables — it was common knowledge among astronomers working in that epoch. Since 2004, however, the situation has been dramatically changing thanks to the increased ac-

tivity of sky surveys and amateur astronomers, and the number of WZ Sge-type dwarf novae discovered in a year reached nearly ten in 2014. Since much information has been collected since the last compilation of WZ Sge-type dwarf novae (Kato et al. 2001) and collection of early superhumps in various objects (Kato 2002), we are tempted to review the recent progress of this field.

In this paper, we mainly focus on optical (and sometimes near-infrared) photometric observations and deal with the current understanding of the outburst phenomenon, superhumps and early superhumps, evolutionary status and related topics. We do not deal with other interesting topics related to WZ Sge-type objects, such as pulsation of the white dwarf (e.g. Warner, van Zyl 1998; Szkody et al. 2010) and hard X-ray emission during outburst (e.g. Senziani et al. 2008). This paper also does not deal with detailed spectroscopic analysis (including Doppler tomograms), oscillations, ultraviolet and X-ray observations.

The paper is loosely arranged in the order of the definition (and historical background), observations, theories and interpretation. However, the topics are often related each other, we put some theoretical background and short interpretations in the observation part, so that readers can know implications of observational features more easily.

In this paper, we use abbreviated names whose full

names are listed in table 6. The data referred to as Kato et al. (2009a)–Kato et al. (2015) include the public data from the AAVSO International Database¹.

2. Definition

2.1. Historical Development

When the class “WZ Sge-type dwarf novae” was first proposed after the dramatic discovery of the second historical outburst of WZ Sge (see e.g. Patterson et al. 1981), two cataclysmic variables UZ Boo and WX Cet were reported to comprise a distinct sub-group of the dwarf novae together with WZ Sge (Bailey 1979). The properties of this subgroup were that they show (b1) large outburst amplitudes (approximately 8 mag), (b2) slow declines from outbursts, and (b3) long intervals between outbursts. Among these properties, (b1) and (b3) have been generally used to define this class. The property (b2) probably referred to the long duration of the outbursts (now confirmed to be superoutbursts) rather than the decline rate itself. Observation of WX Cet in quiescence by Downes, Margon (1981) supported the spectroscopic similarity of WX Cet to WZ Sge proposed in Bailey (1979).

Vogt, Bateson (1982) used in their famous atlas of southern and equatorial dwarf novae a classification of WZ Sge-type dwarf novae. This publication listed WX Cet, AL Com, UZ Boo, V890 Aql and WZ Sge as WZ Sge-type objects and RZ Leo, GO Com, V551 Sgr, YY Tel, AO Oct and VY Aqr as candidate WZ Sge-type objects. This list was selected by the low outburst frequency and large outburst amplitudes, and some objects were included simply due to the lack of observations or overestimation of the amplitudes (among them, V890 Aql turned out to be a chance mis-identification of an asteroid: Samus 1982a; Samus 1982b). Richter (1986b) also suggested V358 Lyr to be a candidate WZ Sge-type dwarf nova. Richter (1986a) provided a list of objects which were considered to be WZ Sge-type dwarf novae. This list consisted of VY Aqr, UZ Boo, WX Cet, AL Com, DO Dra, V592 Her, RZ Leo, WZ Sge and BZ UMa and was apparently selected by the long (more than 4 yr) recurrence time. Mukai et al. (1990) listed BC UMa as an object having properties similar to WZ Sge.

Downes (1990) provided a slightly modified set of properties: (d1) The duration of the outbursts is greater (weeks, versus days in ordinary dwarf novae), (d2) the time between outbursts is greater (years versus months), and (d3) the size of the outbursts is greater (6–8 mag vs 2–5 mag). Downes (1990) also listed a possible property (d4) the strengths of their emission lines in quiescence are greater than those of typical dwarf novae.

O’Donoghue et al. (1991) studied WX Cet in detail. After a comparison of objects suggested to be similar to WZ Sge, O’Donoghue et al. (1991) concluded that there is no reason to retain the distinction between WZ Sge and SU UMa subclasses of dwarf novae. The conclusion

was mainly drawn from the presence of normal (short) outbursts and superhumps in what were supposed to be WZ Sge-type objects — these properties were then considered as properties common to SU UMa-type dwarf novae. O’Donoghue et al. (1991), however, summarized the objects listed in Vogt, Bateson (1982), Richter (1986a), Downes (1990) and Mukai et al. (1990) and provided a convenient list of “properties of possible, probable and certain WZ Sge stars”, which has, rather ironically, become the prototypical collection of WZ Sge-type candidates.

It would be noteworthy that these “classical” definitions did not include the lack (or rarity) of short outbursts, which has long been discussed particularly in WZ Sge.

2.2. Tremendous Outburst Amplitude Dwarf Novae

In the late 1980s, observations of faint CVs with CCDs became more popular (e.g. Howell, Jacoby 1986). Howell’s group systematically observed previously neglected faint CVs in high Galactic latitudes and found many short-period systems (Howell et al. 1987; Howell, Szkody 1988; Howell et al. 1988; Szkody et al. 1989; Howell, Szkody 1990; Howell et al. 1990). During the course of these surveys, Howell, Szkody (1990) noticed that (for systems below the period gap) the mean outburst amplitude for the halo dwarf novae is 3 mag greater than for the disk dwarf novae. They finally reached a concept of “tremendous outburst amplitude dwarf novae” or TOADs (Howell et al. 1995a). The term TOADs was sometime used as a synonym of WZ Sge-type dwarf novae.

Both the historical definitions (subsection 2.1) and TOADs, however, suffered from strong contamination of ordinary SU UMa-type dwarf novae, since they were primarily based on the amplitude of outbursts (especially in TOADs, the amplitude was the sole criterion) and the second proposed prototype, WX Cet, was, after all, shown to be a rather normal SU UMa-type dwarf nova (O’Donoghue et al. 1991; Rogoziecki, Schwarzenberg-Czerny 2001; Kato et al. 2001a) whose historical outbursts were missed likely due to poor observational conditions. There were many similar cases, such as VY Aqr (della Valle, Augusteijn 1990; Augusteijn 1994; Patterson et al. 1993).

Such a contamination of ordinary SU UMa-type dwarf novae to the proposed subclass unavoidably blurred the border, if any, between SU UMa-type and WZ Sge-type subclasses and complicated the discussion. It was a natural consequence that the distinction between WZ Sge and SU UMa subclasses could not be confirmed (O’Donoghue et al. 1991). O’Donoghue et al. (1991) even stated that the extreme behavior of the WZ Sge stars is not a result of the approach of the secondary towards the limit of its evolution as a non-degenerate star at an orbital period (P_{orb}) of ~ 80 min, perfectly contrary to the current understanding. This conclusion was probably a result of inclusion of DO Dra, which is an outbursting intermediate polar with long cycle lengths (Patterson et al. 1992; Wenzel 1983); the contamination by various classes of objects apparently seriously damaged the discussion around that time.

The case for TOADs was even more serious, since they were classified only by the amplitude of outbursts, and

¹ <<http://www.aavso.org/data-download>>.

Patterson et al. (1996) severely criticized the distinction of dwarf novae by outburst amplitudes by showing the continuous distribution of outburst amplitudes.

2.3. *Early Superhumps*

In the meantime, several objects which had been proposed to be similar to WZ Sge-type dwarf novae underwent a dramatic outburst. This epoch coincided with the second-stage progress of CCD photometry in CV research: using small telescope(s) and CCD to detect short-term variations during outburst, which was first systematically conducted by the author (e.g. Kato et al. 2004c).

The first case was in HV Vir in 1992 April (Schmeer et al. 1992). During the early stage of the outburst², double-wave periodic modulations were detected. Although these modulations were detected independently by different groups (Barwig et al. 1992; Mendelson et al. 1992; Leibowitz et al. 1994 and our group), the distinction of these modulations from (ordinary) superhumps should await yet another object. Our result of HV Vir in 1992 was published in Kato et al. (2001) following modern interpretation.

The second case was in AL Com in 1995 [although there were outbursts of LL And in 1993 December (Kato 2004; Howell, Hurst 1996) and UZ Boo in 1994 August (cf. Kato et al. 2001), the observational condition for these objects was not favorable enough to securely characterize the nature of these objects]. The 1995 outburst of AL Com was well-observed and the existence of double-wave modulations during the early stage of the outburst was established (Kato et al. 1996; Patterson et al. 1996; Howell et al. 1996; Nogami et al. 1997). Patterson et al. (1996) suggested that their period is almost same as the orbital period.

These double-wave modulations having period equal to the orbital period were also recorded in the 1978–1979 outburst of WZ Sge (Patterson et al. 1981), and they were started to be recognized as properties unique to WZ Sge-type dwarf novae (cf. Matsumoto et al. 1998). These modulations were called early superhumps (Kato et al. 1996), outburst orbital humps (Patterson et al. 1996) or superorbital modulations (Howell et al. 1996). Since the term “superorbital period” is now widely used in different meaning in X-ray binaries (e.g. Ogilvie, Dubus 2001), the last name is rather confusing and is not used for these double-wave modulations during outburst. Osaki, Meyer (2002) proposed to use “early humps”. Now the term “early superhumps” appears to be used most frequently and we use this term in this paper.

The common existence of early superhumps almost exclusively in dwarf novae with infrequent large-amplitude, long-lasting outbursts became clearer as new observations became available. Although early observations are still somewhat less clear (EG Cnc in 1996 November–December: Matsumoto et al. 1998; Patterson et al. 1998;

Kato et al. 2004a), RZ Leo in 2000 December (Ishioka et al. 2001), two dramatic outbursts of WZ Sge in 2001 July and AL Com in 2001 May (Ishioka et al. 2002; Patterson et al. 2002; Kato et al. 2009a) led to secure and impressive detection of these modulations. After these detections, early superhumps have been regularly detected in dwarf novae with similar systems and the existence of early superhumps gradually became the defining characteristics of WZ Sge-type dwarf novae.

This classification received support from theoretical consideration. Osaki, Meyer (2002) identified early superhumps as manifestation of the 2:1 resonance [note that Lin, Papaloizou (1979) was the first to point out that double peaked light curve in WZ Sge can be related to the 2:1 resonance]. This resonance is almost impossible to achieve for ordinary SU UMa-type dwarf novae, and only objects with extreme mass-ratios are expected to show early superhumps. The modern definition of WZ Sge-type dwarf novae showing early superhumps as manifestation of the 2:1 resonance is favorable in several respects: (1) it is based on the physical mechanism involved in variation, (2) the double-wave profile of early superhumps is very characteristic and they can be identified even if the orbital period is not known, and (3) the objects showing these early superhumps comfortably fit the classical definition of WZ Sge-type dwarf novae. As discussed later, the appearance of early superhumps is inclination-dependent and it is problematic that not all objects achieving the 2:1 resonance show early superhumps. Although it is widely accepted that objects showing early superhumps (persisting at least several days) are classified as WZ Sge-type dwarf novae, the classification is somewhat ambiguous for objects without detectable early superhumps.

2.4. *Rebrightenings*

Since the early period on, the complexity of outburst light curves of WZ Sge-type dwarf novae (and candidates) received attention. A short (~ 1 d) dip was noticed in AL Com as early as Bertola (1964). Ortolani et al. (1980) also noticed a similar dip in WZ Sge during the 1978–1979 outburst of WZ Sge. Richter (1982) stated that dips in the declining branches of the outbursts might be a distinctive feature of WZ Sge-type stars (cf. Richter 1992). Duerbeck (1987) and Howell, Szkody (1988) also remarked on this phenomenon. A collection of light curves showing a variety of complexity can be seen in Richter (1992).

Although the dip phenomenon was recorded in WZ Sge and AL Com, the phenomenon now referred to as rebrightenings (or echo outbursts) received attention since the detection of two rebrightenings in an X-ray transient V518 Per = GRO J0422+32. Kuulkers et al. (1996) suggested that X-ray transients and WZ Sge-type dwarf novae (referred to as TOADs in their paper) have common characteristics by presenting the two post-outburst rebrightenings in UZ Boo which underwent an outburst in 1994. Although this conclusion was based on visual observations and the existence of multiple rebrightenings in UZ Boo

² Most of outbursts in WZ Sge-type dwarf novae are superoutbursts. In this paper, the outburst of this class actually refers to superoutburst (for easier readability) unless otherwise mentioned.

was somewhat doubtful³, the existence of four rebrightenings was confirmed during the 2003–2004 and 2013 outbursts (Kato et al. 2009a; Kato et al. 2014a).

The epoch-making phenomenon was the outburst of EG Cnc in 1996 November–December. Six rebrightenings were detected by a large collaboration mainly conducted by the VSNET team (Kato et al. 2004c). The phenomenon was announced in real-time on the internet and a theoretical paper Osaki et al. (2001) was issued with a reference to our preprint.

Since then, a long rebrightening similar to those in WZ Sge and AL Com was recorded in CG CMa (Kato et al. 1999). Recent observations have indicated WZ Sge-type dwarf novae are frequently associated with multiple rebrightenings (see e.g. Nakata et al. 2013b), and this phenomenon has been considered to be potential defining characteristics of WZ Sge-type dwarf novae. Some authors classify objects with multiple rebrightenings as WZ Sge-type dwarf novae (Mroz et al. 2013). The physical mechanism of multiple rebrightenings, however, is not as well understood as early superhumps, and it is not known whether ordinary SU UMa-type dwarf novae never show multiple rebrightenings.

2.5. Modern Criteria

In recent years, objects with large-amplitude outbursts (typically ~ 8 mag, at least greater than 6 mag) that exhibit early superhumps at least for several days during the early stage of long outbursts have been unambiguously classified as WZ Sge-type dwarf novae. As theory predicts (Osaki, Meyer 2002), some low-inclination systems do not show detectable early superhumps. In such cases, the existence of a long-duration segment (approximately 10 d or more) without short-term photometric variations before starting to exhibit ordinary superhumps has been usually considered as a signature of “unobservable” early superhumps. Objects showing large-amplitude outbursts with this signature have been usually identified as WZ Sge-type dwarf novae although there is currently no way to confirm that the 2:1 resonance is indeed working. The presence of multiple rebrightenings is considered to be supportive evidence.

These criteria general match the historical category. The shortest measured intervals of long outbursts in objects satisfying these criteria is slightly over 4 yr (EZ Lyn; Pavlenko et al. 2012; Kato et al. 2012a and OT J213806; Kato et al. 2015), and one exceptional case of ~ 450 d AL Com in 2013 and 2015, the latter lacked the stage of early superhump but showed a long rebrightening.

The problem of this criteria will be discussed later.

³ Later examination of the AAVSO observations indicated mixed detections and non-detections of rebrightenings around the same epoch, making it difficult to determine which observations were true detections of rebrightenings.




Fig. 1. Year of recognition of WZ Sge-type properties. The sample is the objects in table 6. Note that the year 2015 includes only January.

3. Statistics

3.1. Discovery Statistics

Figure 1 indicates the discovery statistics of WZ Sge-type objects. The year when the WZ Sge-type nature (in the modern sense) was recognized is used to draw this figure. Up to 2000, WZ Sge-type objects were rare and they were sometimes called by “the n -th WZ Sge-type object”. There has been a dramatic increase of the number since 2004, when ASAS-3 (Pojmański 2002) started discovering new WZ Sge-type objects. In recent years, the tendency is more striking after the increase of discovery by CRTS (Drake et al. 2009), MASTER (Gorbovskoy et al. 2013) and ASAS-SN (Shappee et al. 2014; Davis et al. 2015). In the most recent year of 2014, the contribution of the surveys are: CRTS (1), MASTER (1), ASAS-SN (9) and amateur discoveries (6). Although ASAS-SN has been discovering a great number of WZ Sge-type objects, wide-field survey of bright transients by amateur astronomers still has a great impact in this field.

3.2. Maximum Magnitude

Figure 2 shows the distribution of maximum recorded magnitudes. Note that true maxima were not always recorded. Although the incompleteness becomes more apparent for objects fainter than magnitude 13, the detection is already apparently incomplete even brighter than magnitude 10 since it is well-known that uniformly distributed stars have a number count $\log N(m) = 0.6m + C$, where m is the magnitude and $N(m)$ is the number of stars having apparent magnitudes brighter than m . This relation expects a threefold increase of objects by one magnitude. The data suggest that only half of WZ Sge-type objects having maximum magnitudes 9–10 mag have been discovered.

Just for completeness, we have studied the distribution




Fig. 2. Distribution of maximum recorded magnitudes. The sample is the objects in table 6.

of the months when these objects were recognized. The smallest number is five in May and the largest number is 12 in November. A χ^2 -test yielded a p-value of 0.58, indicating that the distribution cannot be considered as strongly different between different months. This is probably a result of significant contribution of southern observers and modern surveys in the southern hemisphere.

3.3. Outburst Amplitude

Figure 3 shows the distribution of outburst amplitudes. This figure can be compared to figure 13 (distribution of outburst amplitude of all dwarf novae) in Patterson et al. (1996), who criticized the concept of TOADs by indicating that the distribution of amplitudes makes a smooth continuum. Although we no longer use the amplitude as the primary criterion of WZ Sge-type dwarf novae, it is evident that WZ Sge-type dwarf novae occupy the region with largest outburst amplitudes. The lower 75% quantile is 6.9 mag, indicating that the majority of WZ Sge-type dwarf novae have outburst amplitudes larger than 7 mag. The median value is 7.7 mag.

The largest value (9.5 mag) is recorded in SSS J224739. Since the measurement of outburst amplitudes is severely limited for large-amplitude systems and fainter objects, the present statistics is severely biased for objects with smaller amplitudes. Measurements of more reliable quiescent magnitudes to determine the amplitudes are desired for many less-studied objects, although WZ Sge-type dwarf novae usually stay 1 mag or more brighter than the pre-outburst (for example, V455 And has not returned to the pre-outburst level even six years after the 2007 outburst according to the CRTS data), these measurements would require additional years.

The objects with smallest values are V1108 Her, EZ Lyn, EG Cnc, PT And, SDSS J161027 and SS LMi. The quiescent magnitude of V1108 Her is difficult to measure due to the close companion. The magnitudes are approximate for PT And and SS LMi, and may be underestimated (or these objects may be borderline objects). EZ




Fig. 3. Distribution of outburst amplitudes. The sample is the objects in table 6 and objects with lower limit for the amplitude are excluded.

Lyn is an eclipsing system and the outburst amplitude is expected to be smaller than if the object were seen from pole-on. Considering these examples, many of WZ Sge-type objects with low outburst amplitudes in this survey may not reflect the true strength of the outburst. This would strengthen that the majority of WZ Sge-type objects have amplitudes larger than 7 mag. EG Cnc would worth attention. This object showed multiple rebrightenings. While Patterson et al. (1998) suggested this object to be a candidate period bouncer (see subsection 7.8), Nakata et al. (2013b) showed that objects with multiple rebrightenings are not necessarily good candidates for period bouncers. Determination of physical parameters and detailed observations of the next superoutburst of this object are still highly desired.

3.4. Orbital Periods

Figure 4 illustrates the distribution of the orbital periods of WZ Sge-type dwarf novae. For the objects without orbital periods (or periods of early superhumps), we estimated them using the updated relation between the orbital and superhump periods (P_{SH} , equation 6 in Kato et al. 2012a). The WZ Sge-type dwarf novae mostly have orbital period shorter than 0.06 d and comprise the recently identified “period spike” of the CV period distribution (Gänsicke et al. 2009; Kato et al. 2015). The 50% quantiles of the distribution is 0.0553–0.0592 d and the median value is 0.0569 d. There are several outliers, which are either long-period systems (long-period objects having properties common to the short-period WZ Sge-type dwarf novae) or the EI Psc-type object with the evolved secondary core.

3.5. Intervals between Superoutbursts

Figure 5 shows the distribution of intervals between superoutbursts using the data in table 6. If multiple outbursts were recorded in the same object, significantly (more than twice) longer intervals were not used since out-




Fig. 4. Distribution of orbital periods. The sample is the objects in table 6. For the objects without orbital periods (or periods of early superhumps), we estimated them using the updated relation between the orbital and superhump periods (equation 6 in Kato et al. 2012a).




Fig. 5. Distribution of intervals between superoutbursts. The sample is the objects in table 6. If multiple outbursts were recorded in the same object, significantly (more than twice) longer intervals were not used since outbursts may have been missed by chance between these outbursts. Long-period objects and EI Psc-type object are not included.

bursts were likely missed by the lack of observations between these outbursts. Of course, not all outbursts were detected in the objects with multiple known outbursts and these values contain intervals longer than the actual ones. Instead, many objects have only one outburst detections and it is impossible to determine the intervals. If there are recently discovered objects with cycle lengths longer than 10 yr, they are less likely included in this figure. Keeping these restrictions in mind, we can see that the majority of WZ Sge-type objects have shorter recurrence time than the 23–33 yr in WZ Sge. The median value is 11.5 yr. The shortest known interval between well-confirmed superoutbursts in WZ Sge-type objects is 450 d in AL Com (Kimura et al. 2015; not included in this statistics).

3.6. Normal Outbursts

The absence of normal outbursts has frequently been a defining characteristics of WZ Sge-type objects. This is still true for WZ Sge (see, however, a discussion is Patterson et al. 1981; there are still gaps in observation and we cannot exclude a possibility of a short normal outburst). Normal outbursts have been recorded in some objects: AL Com [1974? and 2003 (Kato et al. 2014a)], EG Cnc [2009 (Templeton 2009, Lange 2010)], RZ Leo [1989 (O’Donoghue et al. 1991)], although RZ Leo may be better classified as a long-period system (subsection 7.9). Based on the present statistics, normal outbursts are indeed rare, if not absent, in WZ Sge-type objects.

3.7. Eclipsing Systems

Among nearly 100 WZ Sge-type objects in table 6, only four systems are eclipsing. It is also worth noting only two systems (EZ Lyn and MASTER J005740) were shown to be strongly eclipsing during outbursts [eclipses in WZ Sge were apparent only in certain stages or beat phases of outbursts, cf. Patterson et al. (2002)]. If the orbital planes

are randomly orientated, we can expect 25% objects are eclipsing if objects with inclinations more than 75° are observed as eclipsing systems (the value is from Araujo-Betancor et al. 2005a). The fractions will be 17% and 9% if the inclination limits are 80° and 85° , respectively. The observed eclipsing systems are too few compared to this expectation. This may be a combination of (1) possible selection effect that highly inclined systems are less luminous in outburst, and are less frequently detected as transients or less frequently observed to search for superhumps, and (2) WZ Sge-type systems have very extended accretion disks during the stage of early superhumps, and it is difficult to distinguish the profile of eclipses from that of early superhumps (for numerical model calculations, see Uemura et al. 2012, Kato et al. 2014a). The selection effect (1) is probably a minor contribution since many eclipsing SU UMa-type dwarf novae have been discovered. We can expect observations of these WZ Sge-type dwarf novae in quiescence will identify more eclipsing systems.

4. Phenomenon in Outbursting WZ Sge-Type Dwarf Novae

4.1. Outburst Morphology and Rebrightenings

The initial part of the superoutburst of WZ Sge-type dwarf novae usually has a steeper decline (this period approximately corresponds to the period with early superhumps), and this part is the viscous depletion period (cf. Osaki 1995a; Osaki, Meyer 2003) which has a power-law type (faster than exponential) decay (Cannizzo et al. 1990; Cannizzo 1996). After this phase, the outburst enters the exponential decline phase (slow decline; subsection 4.5). This part of the light curve is essentially the same as those of ordinary SU UMa-type dwarf novae. WZ Sge-type dwarf novae, however, show a variety of post-outburst rebrightenings or “dips” in the light curve as

already introduced in subsection 2.4.

Imada et al. (2006) was the first to classify the morphology of rebrightenings. The four classes are: type-A outbursts (long-duration rebrightening), type-B outbursts (multiple rebrightenings), type-C outbursts (single rebrightening) and type-D outbursts (no rebrightening). Originally, type-A outburst was introduced due to the similarity of light curves between WZ Sge and AL Com. A more closer examination, however, indicated that the 2001 outburst of WZ Sge is composed of low-amplitude multiple rebrightenings (Patterson et al. 2002; Osaki, Meyer 2003; Kato et al. 2009a), and it looks like that type-A and type-B form a smooth continuum (Meyer, Meyer-Hofmeister 2015). For this reason, we write type-A/B for outbursts with a long-duration rebrightening composed of low-amplitude multiple rebrightenings. There are, however, long-duration rebrightenings without detectable low-amplitude multiple rebrightenings (such as AL Com in 2013), and we refer them as type-A. Modern examples of “textbook” light curves are shown in figures 6, 7, 8. For type-B (and type-A/B) outbursts, Meyer, Meyer-Hofmeister (2015) provided an excellent summary of light curves and relationship between the amplitudes and intervals of rebrightenings.

Type-E was introduced by Kato et al. (2014b) after the detection of two objects with double superoutbursts (SSS J122221: Kato et al. 2013b; OT J184228: Kato et al. 2013a). An example is shown in figure 9. These objects are considered to be good candidates for period bouncers (Kato et al. 2013b).

4.2. Reproducibility of Rebrightening

It has not been well investigated whether the same star shows or tends to show the same type of rebrightenings. Although a comparison of the 1913, 1946 and 1978–1979 outbursts of WZ Sge seemed to show noticeable difference between outbursts (Patterson et al. 1981), this study was before the recognition of rebrightening phenomenon and the presentation of the data may have been biased. The AAVSO page⁴ described “this may be due to the lack of data for the 1946 outburst since the decline and recovery is fast, if there is no continuous data it would be easy to miss” and the seeming difference between different outbursts may have been caused by lack of observations.

In other objects, Uemura et al. (2008b) observed the rebrightening part of AL Com and reported the rebrightening in 2007–2008 was composed of discrete short outbursts in contrast to the 1995 and 2001 ones. The most recent comparison of different outbursts in AL Com suggests that the rebrightening in this object tends to be reproducible, and the rebrightening in 2007–2008 was composed of small brightenings with amplitudes less than 1 mag, which is not different from the type-A/B rebrightening in WZ Sge in 2001 (Kato et al. 2014a). AL Com underwent an unusually faint superoutburst in 2015, but showed type-A rebrightening (Kimura et al. 2015).

Objects with discrete short outbursts (type-B rebright-




Fig. 6. Upper: 2013 superoutburst of AL Com (type-A rebrightening). The data are taken from Kato et al. (2014a). The data points were binned to 0.019 d for the superoutburst and to 1 d after the superoutburst. Lower: 2001 superoutburst of WZ Sge (type-A/B rebrightening). The data are taken from Kato et al. (2009a). The data points were binned to 0.019 d for the superoutburst and to 1 d after the superoutburst.

enings) tend to show the same type of rebrightenings in the cases when multiple outbursts were recorded, although the number of such objects has been still small. The numbers of rebrightenings, however, can vary. These objects are UZ Boo (1993, 2004: Kuulkers et al. 1996, Kato et al. 2009a), EZ Lyn (2006, 2010: Pavlenko et al. 2012; Kato et al. 2012a).

In OT J213806, with type-D outbursts, remarkably different features (particularly the duration of the plateau phase) were observed between the 2010 and 2014 outbursts (Kato et al. 2015). Neither outburst, however, showed a rebrightening.

4.3. Case Study of WZ Sagittae

We examine here the historical outbursts of WZ Sge. The observations for 1913 and 1946 shown in Patterson et al. (1981) were from observations in Mayall (1946) rather than from the AAVSO database as referred to in

⁴ <http://www.aavso.org/vsots_wzsge>.




Fig. 7. Upper: 2010 superoutburst of EZ Lyn (type-B rebrightening). The data are taken from Kato et al. (2012a). The data points were binned to 0.019 d before BJD 2455509 and to 1 d after this. Lower: 2012 superoutburst of MASTER J211258 (type-B rebrightening). The data are taken from Nakata et al. (2013b). The data points were binned to 0.019 d.




Fig. 8. Upper: 2004 superoutburst of FL Psc (type-C rebrightening). The data are taken from Kato et al. (2009a). The data points were binned to 0.019 d. Lower: 2007 superoutburst of GW Lib (type-D rebrightening). The data are taken from Kato et al. (2009a). The data points were binned to 0.019 d.

the paper.

We first examined the magnitude scale since there was a possibility of systematic difference from the modern scale (cf. footnote 4 of Kato et al. 2001). The magnitudes of photographic comparison stars listed in Mayall (1946) have been found to agree to Tycho-2 B magnitudes or CCD B magnitudes for fainter stars within 0.2 mag. It has been confirmed that the 1913 and 1946 observations recorded the object in a system equivalent to the modern B band. AAVSO observations of the 2001 outburst suggest that WZ Sge had $B - V = -0.1$ and $U - B = -1.0$ 1 d after the maximum. If the 1913 and 1946 photographic plates correctly reproduced the modern B -band, the recorded maxima were brighter than the later ones in 1978 and 2001. If the plates had sensitivity to the U light, the recorded brighter magnitudes in 1913 and 1946 may have been attributed to the sensitivity, and the outburst amplitude may be overestimated of we treat the magnitudes of the 1913 and 1946 outbursts as V magnitudes.

We examined the light curve of the rebrightening part.




Fig. 9. 2011 superoutburst of OT J184228 (type-E rebrightening). The data are taken from Kato et al. (2013a). The data points were binned to 0.019 d.

Since the original magnitudes were not available in publication, we have extracted the values from the figures in Mayall (1946). The errors of dates and magnitudes were expected to be less than 1 d and 0.1 mag, respectively, and will not affect the following discussion. We also used the AAVSO database for visual observations of the 1946 outburst.

The comparison is shown in figure 10. In this figure, the start of the 2001 superoutburst was artificially shifted by 4 d. This measure was based on recent two examples (OT J213806: Kato et al. 2015 and AL Com: Kimura et al. 2015) in which superoutbursts of different extent were observed and the difference between superoutbursts has been found in the duration of the early part of the superoutburst. The result indicates that the light curves of the 1978 and 2001 superoutbursts are very similar with a fluctuating long rebrightening, except that the initial part of the 2001 is shorter. This characteristic is in good agreement with OT J213806 in 2010 and 2014. Although the light curve of the 1946 in Patterson et al. (1981) looks like to show the absence of rebrightening(s), we should note that there was a gap of observations for 8 d in Mayall (1946). Two AAVSO observations during this gap showed brighter (12.0–12.5) magnitudes. Although these visual observations at faint magnitudes may have not been very reliable, we cannot rule out the possibility of a rebrightening during this gap. A long rebrightening can be safely excluded. The 1913 superoutburst was not very densely observed. Since the other three superoutbursts experienced rapid fading 24–29 d after the peak brightness, we consider it highly likely that the object faded after the second final observation at 26 d. If this is the case, the final observation was obtained during a rebrightening.

In summary, although there was strong evidence against the presence of a long rebrightening in the 1946 superoutburst, the other three superoutburst (certain for the 1978 and 2001 ones) showed similar rebrightenings.

4.4. Properties of Rebrightenings

The basic properties of repetitive rebrightenings are summarized in Meyer, Meyer-Hofmeister (2015). Since Meyer, Meyer-Hofmeister (2015) only dealt with multiple rebrightenings (our type B or A/B), we deal with them first. We provide an updated table in table 1. This table only includes type-B objects in our category and does not include type-A/B and “mini-rebrightenings” in the SU UMa-type object V585 Lyr (not classified as a WZ Sge-type object in this paper). For the objects listed in this table, the values in Meyer, Meyer-Hofmeister (2015) were included without modification. For references of other objects, see table 6.

The number of rebrightening ranges from 2 to 11 (12, including WZ Sge as in Meyer, Meyer-Hofmeister 2015). The amplitude is positively correlated with intervals (see figure 5 in Meyer, Meyer-Hofmeister 2015). Our new object 1RXS J023238 gives a support, while OGLE-GD-DN-014 shows a smaller amplitude. It may be that OGLE-GD-DN-014 may belong to a population different from short-period WZ Sge-type objects (see a comment in ta-




Fig. 10. Comparison of superoutbursts of WZ Sge. The data were binned to 1 d and shifted in magnitude. The dashed lines are added when continuous data are available to aid recognizing the variation. The data for the 2001 superoutburst were from our CCD data (Ishioka et al. 2002). The data for the 1978 superoutburst were from AAVSO observations. The data for the 1946 superoutburst were from Mayall (1946) and AAVSO observations. The data for the 1913 superoutburst were from Mayall (1946). The start of the 2001 superoutburst was artificially shifted by 4 d so that the time of the “dip” generally agree to the others.

Table 1. WZ Sge-type objects with multiple rebrightenings

Object	Year	N_{reb}^*	t_{reb}^\dagger [d]	Δm [mag]
UZ Boo	1994	≥ 2	–	–
UZ Boo	2003	4	3.6	2.4:
UZ Boo	2013	4	4.0	3.0
DY CMi	2008	6	4.7	2.9
EG Cnc	1996	6	7.3	3.3
VX For	2009	5	4.5	2.9
QZ Lib	2004	4	5.5	2.8
EZ Lyn	2006	11	2.7	1.6
EZ Lyn	2010	6	4.9	2.5
EL UMa	2010	≥ 5	4.0	2.2
1RXS J023238	2007	≥ 4	9.2	3.7
ASASSN-14cv	2014	8	4.1	2.7
MASTER J085854	2015	2	4.9	2.5
MASTER J211258	2012	8	4.7	2.6
PNV J171442	2014	5	4.3	1.9
TCP J233822	2013	2	4.9	2.0
OGLE-GD-DN-001	2007	4	5.0	2.8
OGLE-GD-DN-014	2006	2	10.1	2.5

*Number of rebrightenings.

† Average interval of rebrightenings.

‡ Average amplitude of rebrightenings.

ble 6).

Meyer, Meyer-Hofmeister (2015) also noted basic properties of multiple rebrightenings: (1) the minimum luminosity of the rebrightenings is higher than the quiescence level, (2) the maximum luminosity is on a smooth extension of the decline from the main superoutburst and the minimum luminosity decreases in parallel to the maximum luminosity (somewhat rephrased and supplemented by our own interpretation of the data), (3) (they usually show) more rapid brightness increase and the slower brightness decrease.

These properties are mostly general to rebrightenings. There are, however, some exceptional cases. In EG Cnc, the initial five rebrightenings had rapid brightness increase, suggesting outside-in type outbursts, while the final one had slower brightness increase, suggesting an inside-out type outburst (Kato et al. 2004a). Following the first rebrightening of EG Cnc, there was a small and slowly rising rebrightening, which did not reach the brightness of other rebrightenings (Kato et al. 2004a). In EG Cnc, the interval of the final rebrightening was longer than the others, and a similar trend was seen in the final rebrightening of EZ Lyn in 2010 (figure 35 in Kato et al. 2012a) the type-A/B object WZ Sge (figure 36 in Kato et al. 2012a).

In well-observed systems, superhumps are continuously seen during the rebrightenings. The amplitudes of superhumps are inversely correlated with the system brightness (e.g. Patterson et al. 1998; Patterson et al. 2002), and the pulsed flux of the superhumps was almost constant despite large variation due to rebrightenings. There was some evidence that the flux of the superhumps decreases before the termination of the rebrightening phase (figures 35, 36 in Kato et al. 2012a).

In cases of long rebrightenings without (type-A outburst), the long rebrightening is often associated by a precursor outburst and superhumps look like to appear and evolve again during the rebrightening [AL Com: Nogami et al. (1997), Kimura et al. (2015); CSS J174033: T. Ohshima et al. in preparation; ASASSN-13ax and ASASSN-13ck showed a deep fading after the initial rise (Kato et al. 2014b)]. The rising phase of the rebrightening in the 1995 outburst of AL Com was slow, suggesting that it was an inside-out type outburst Nogami et al. (1997).

We found “mini-rebrightenings” between the main superoutburst and the single rebrightening in the Kepler data of V585 Lyr (ordinary SU UMa-type dwarf nova) (Kato, Osaki 2013a). Although this phenomenon was reproduced in the second recorded superoutburst in Kepler data, we still do not have corresponding data in other objects in the ground-based observations. It is not known whether the same phenomenon is present in WZ Sge-type objects.

4.5. Slow Fading Rate

Cannizzo et al. (2010) reported that the fading rates of WZ Sge is much faster than the Kepler objects, V344 Lyr and V1504 Cyg, and suggested that this different could arise from the strong dependence of the viscosity in quies-

cence. Cannizzo et al. (2010) interpreted that smaller viscosity in quiescence gives rise to a larger surface density at the start of the superoutburst and hence a steeper viscous decay. His analysis, however, used different segments between ordinary SU UMa-type dwarf novae (linearly fading part) and WZ Sge (initial rapid fading). When restricted to linearly fading part, Kato et al. (2014b) found no difference of fading rates between ordinary SU UMa-type dwarf novae and WZ Sge-type dwarf novae. Kato et al. (2014b) found that the fading rate follows the theoretically expected dependence $P_{\text{orb}}^{1/4}$.

Kato et al. (2014b) also found that some WZ Sge-type dwarf novae tend to show significantly slower fading rates, and attributed this deviation to smaller disk viscosity in the hot state. Kato et al. (2014b) considered that this viscosity reflects the tidal strength and suggested that objects with slower fading rates have smaller mass ratios ($q = M_2/M_1$) and they are good candidates for period bouncers (subsection 7.8).

4.6. Global Color Variation

There have been a number of studies of global color variations of WZ Sge-type dwarf novae. Patterson et al. (1998) observed EG Cnc in 1996–1997 and obtained $B - V$ and $V - I$ close to zero during the superoutburst plateau. The object became redder between the rebrightenings, in particular $V - I$ reached 0.7. Howell et al. (2004) also noted the similar trend in WZ Sge in 2001. The object was bluest ($B - V \sim -0.1$) around the brightness peak and became redder ($B - V \sim 0.3$) during the fadings between the rebrightenings. $V - I$ also became red (~ 0.6), but became bluer (~ 0.3) after the end of the rebrightening episodes despite that the object further faded. The $U - B$ values were mostly strongly negative (~ -1.0) except during the rebrightening episodes. The blue color in $U - B$ is similar to other dwarf novae in outburst and quiescence, which is a result of the weak Balmer jump of an outbursting accretion disk and also contribution of strong emission lines in quiescence.

Uemura et al. (2008a) conducted multicolor infrared observations of IK Leo and found an excess in K_s band during the rebrightening phase. Uemura et al. (2008a) considered this excess arises from an optically thin region that is located outside the optically thick disk. Matsui et al. (2009) reported optical and near-infrared color variations of V455 And. Although V455 And showed no rebrightening, the $V - J$ colors remained very red (~ 0.8) at least for 34 d in the post-superoutburst state. Uemura et al. (2008a) modeled the colors by a combination of blackbody and free-free emission, and concluded that the blackbody emission remained at a moderately high temperature (~ 8000 K) for 10–20 d after the superoutburst, suggesting the existence of a substantial amount of gas remaining in the disk. Uemura et al. (2008a) discussed the possibility of such gas as an origin of rebrightenings.

Chochol et al. (2012) also reported red colors for an interval of ~ 10 d after the superoutburst of OT J213806. Nakagawa et al. (2013) reported red colors during the

dip before the rebrightenings in OT J012059. Golysheva, Shugarov (2014) reported red colors (particularly in the I band) for PNV J191501, which did not show rebrightenings. Isogai et al. (2015) studied EZ Lyn in two bands (g' and i') and noted red colors near minimum light of multiple rebrightenings.

From these observations, it has been established that WZ Sge-type dwarf novae show red colors during the rebrightening phase (for objects with multiple rebrightenings) and that the objects without rebrightenings tend to show a prolonged phase of red colors. These observations give a clue in understanding the mechanism of rebrightenings (see subsection 8.3).

4.7. Global Spectral Variation

In this subsection, we concentrate on optical low-resolution spectroscopy and discuss the global variation of the spectra.

Nogami, Iijima (2004) is still the best reference for systematic study of spectral variation of the WZ Sge-type dwarf nova (WZ Sge itself). During the early stage of the outburst (corresponding to the phase of early superhumps, see section 5), the object showed a hot continuum with broad Balmer absorption lines, which are characteristic to dwarf novae in outburst and reflect an optically thick hot accretion disk.

In addition to these features, HeII and the Bowen complex (CIII/NIII) were seen in emission. Although this feature is known to appear in other dwarf novae in some phase (e.g. Hessman et al. 1984) and in high-inclination systems (e.g. Steeghs et al. 1997; Wu et al. 2001), these high excitation lines are most notably observed in WZ Sge-type dwarf novae (particularly during the early stage): GW Lib and V455 And (Nogami et al. 2009), V592 Her (Mennickent et al. 2002), CRTS J090239 (Djorgovski et al. 2008), OT J111217 (vsnet-alert 9782, Kato et al. 2009a), V572 And (Quimby et al. 2005). PNV J191501 (vsnet-alert 15779, Kato et al. 2014a), ASASSN-14cl (Teyssier 2014), PNV J172929 (vsnet-alert 17327, Kato et al. 2015). A less striking cases was V355 UMa (vsnet-alert 12822, Kato et al. 2012a). In some cases, higher excitation lines CIV and NIV have been detected (WZ Sge: Nogami, Iijima 2004), OT J111217: vsnet-alert 9782, Kato et al. 2009a), which have not yet been detected in other types of dwarf novae.

Subsequent development of the spectra were not very striking (Nogami, Iijima 2004): Balmer emission lines gradually turn to emission lines as the system fades.

The most notable feature in the spectra of WZ Sge-type dwarf novae is the presence of Na D absorption, which was first detected in EG Cnc during the rebrightening phase (Patterson et al. 1998). WZ Sge also showed this feature, but was seen during the early stage of the outburst (Nogami, Iijima 2004). The origin of Na D absorption may be different between the rebrightening phase in EG Cnc and outburst phase in WZ Sge. Combined with the red color in the rebrightening phase (subsection 4.6), this Na D absorption gives additional support to the existence of a cool component in the disk.

Since this paper does not intend to give full details of the spectral evolution, please refer to Nogami, Iijima (2004) for detailed spectral development and past observations.

5. Early Superhumps

5.1. General properties

As introduced in subsection 2.3, the most prominent (and likely most discriminative) feature discovered in WZ Sge-type dwarf novae is early superhumps. These modulations are double-wave modulations seen during the initial stage of the superoutburst and have periods extremely close to the orbital period to an accuracy of 0.1% (Ishioaka et al. 2002; Kato et al. 2014a). The currently most promising interpretation is the spiral structure excited by the 2:1 resonance (Osaki, Meyer 2002) and the variation is caused by a geometrical effect. There was a discussion whether the light source is an illumination of the azimuthally extended disk (Kato 2002). Osaki (2003) criticized this interpretation. Maehara et al. (2007) was the first to successfully model the light curve of early superhumps by assuming the azimuthally extended disk with spiral arms. More recently, Uemura et al. (2012) succeeded in mapping the height of the accretion disk of V455 And by using the multicolor light curve of early superhumps by considering the geometrical projection effect and self-eclipse. Uemura et al. (2012) also estimated the illumination effect and concluded that it is a minor one. The model, however, has not been able to reproduce the Doppler tomogram or line variations of V455 And (Uemura et al. 2015) and it would require a new interpretation.

All these modern works suggest that early superhumps can be only seen in high-inclination systems, as modeled by Uemura et al. (2012). Observations also support this interpretation: there was no strong early superhumps in GW Lib (Kato et al. 2009a) which is reported to have a low inclination of 11° (Thorstensen et al. 2002) and all systems with eclipses have high-amplitude early superhumps [e.g. Kato et al. (2009a) for WZ Sge; V455 And and MASTER J005740 Kato et al. (2014a)]. There are intermediate cases such as AL Com which shows prominent orbital variations in quiescence (Patterson et al. 1996; Szkody 1987).

Since figure 1 of Kato (2002) has been frequently used as a “catalog of early superhumps”, we provided an updated figure with much improved statistics (figure 11). We have restricted the objects to those with good statistics, and some objects in Kato (2002) are omitted from this figure. Except EZ Lyn, all object show double-wave modulations having the secondary (around phase 0.5–0.6) maximum brighter. The dip between two maxima (around phase 0.3) is somewhat dependent on the objects, and some objects (like V455 And) only show a shallow dip and the overall profile resembles a “saw-tooth” with a slower rise. Such “saw-tooth”-like profile is usually seen in objects with high amplitudes of early superhumps. A recent example is ASASSN-15hd (vsnet-alert 18552, 18555). In this case, early superhumps were initially almost singly peaked (with a slow rise in contrary to ordinary superhumps) but




Fig. 11. Mean profiles of early superhumps. For the initial five objects, the zero phase corresponds to eclipses. The zero phases of other objects are chosen so that the stronger hump is located around phase 0.6. The year is added after the object name if multiple outbursts have been recorded.

later they became double-humped (vsnet-alert 18573).

5.2. Evolution of Early Superhumps

In well-observed systems, early superhumps grow as the object rises to the outburst maximum (WZ Sge and AL Com: Ishioka et al. 2002). During the rising phase of V455 And (2007), the object did not show any evidence of early superhumps from ~ 6 mag to ~ 1.5 mag before the maximum then these superhumps started to grow quickly (Maehara et al. 2009; figure 12). In WZ Sge (2001), early superhumps became evident around ~ 1.0 mag below maximum (figure 13; Ishioka et al. 2002). These two systems are best studied ones during the growing stage of early

superhumps.

5.3. Amplitude Variation

In well-observed systems, it has been demonstrated that the amplitudes of early superhumps decrease with time. The best example may be figure 2 in Patterson et al. (2002). Note that this figure express the amplitudes in intensity. Since the mean brightness of the object decreases with time, the amplitudes expressed in magnitudes do not decrease so dramatically. In order to illustrate this effect, we provide figure 14, in which both amplitude and intensity variations are given for the entire interval when early superhumps were present. It is now evident that the decrease in amplitude is not so dramatic (particularly in the later phase) as the impression from figure 2 in Patterson et al. (2002).

5.4. Amplitude Statistics

We made a survey of amplitudes of early superhumps (table 2). This table includes the objects which were sufficiently observed to tell the presence of early superhumps. Although amplitudes of early superhumps systematically decrease with time (subsection 5.3), most of objects were not sufficiently observed to follow the variation of amplitudes. We therefore used mean amplitudes for the entire interval when early superhumps were present. In some well-observed objects, amplitudes around the peak brightness are given. The distribution of mean amplitudes of early superhumps indicates that the majority of systems have amplitudes less than 0.05 mag, although there are a small number of objects showing large-amplitude early superhumps up to 0.35 mag. The numbers of the objects having amplitudes of early superhumps larger than 0.02 (0.05) mag are 33 (11) out of a total number of 52. If we can typically detect 0.02-mag early superhumps, we can classify 63% of WZ Sge-type dwarf novae by the presence of early superhumps. If we can detect 0.01-mag early superhumps, this fraction becomes 79%, making this criterion as WZ Sge-type objects more promising than may have been thought.

5.5. Comparison of Amplitude Statistics with Model

In recent works, there has apparently been a consensus that early superhumps are a result of the 2:1 resonance (Osaki, Meyer 2002; Kato 2002; Osaki, Meyer 2003) although this phenomenon was first recognized in history as an enhanced orbital humps (Patterson 1980). Although the nomenclature and the presentation of the figure in Patterson et al. (2002) would give an impression of an enhanced hot spot (as explained in Osaki, Meyer 2002), we should note that Patterson et al. (2002) wrote “a model of this type seems very attractive” and “Osaki, Meyer (2002) provide a lucid explanation for the one prominent feature not previously explained”.

We studied whether the statistics in subsection 5.4 can be explained by an inclination effect. We used a code described in Uemura et al. (2012). We assumed the disk structure reconstructed from observation of V455 And and changed the orbital plane randomly, and examined the




Fig. 12. Evolution of early superhumps in V455 And (2007) The data are from Kato et al. (2009a). The data points were binned to 0.0005 d. Upper: light curve. Lower: residual magnitudes.




Fig. 15. Distribution of mean amplitudes of early superhumps. The data are taken from table 2. Although the majority of systems have amplitudes less than 0.05 mag, a small number of objects show large-amplitude early superhumps up to 0.35 mag.

distribution of expected amplitudes. A result of 100000 trials is shown in figure 16. The result above 0.4 mag is not real, since this model does not treat the geometrical structure of the edge of the disk properly and gives a senseless result for inclinations larger than 82° . We used a constant value for 82° for systems with higher inclinations. The result seems to reproduce the high number of systems with low amplitudes of early superhumps. This result seems to support the geometrical origin of early superhumps. The model, however, predicts a large fraction ($\sim 20\%$) of objects with amplitudes larger than 0.2 mag, which is different from observations (6%). Since the model is rough and the disk model of V455 And may not represent the true disk, this discrepancy may not be a strong contradiction. There may be a possibility that very high-inclination WZ Sge-type systems do not show strong early superhumps.

5.6. Colors

Color variations of early superhumps have been studied in order to constrain the mechanism to produce them. Matsui et al. (2009) was the first to systematically study color variations of early superhumps using (nearly) simultaneous multi-color time-resolved photometry including




Fig. 13. Evolution of early superhumps in WZ Sge (2001) The data are from Kato et al. (2009a) (same as Ishioka et al. 2002). The data points were binned to 0.0005 d. Upper: light curve. Lower: residual magnitudes.

Expected Amplitude of Early Superhumps




Fig. 16. Expected distribution of amplitudes of early superhumps. The amplitudes have been estimated from 100000 randomly oriented systems having the same disk parameter as in V455 And and using the model in Uemura et al. (2012).

infrared bands. In contrast to well-established color variation of superhumps of ordinary SU UMa-type dwarf novae (e.g. Schoembs, Vogt 1980; Hassall 1985; Naylor et al. 1987; van Amerongen et al. 1987; van der Woerd, van Paradijs 1987), early superhumps were found to be redder when brighter. This indicates that the light source of early superhumps is cooler than the underlying component and Matsui et al. (2009) suggested that early superhumps are produced in a vertically extended low-temperature zone at the outermost part of the disk. The reconstruction of the disk geometry by using multicolor photometry of early superhumps by Uemura et al. (2012) was an immediate result of this work. Isogai et al. (2015) performed two-color simultaneous photometry of EZ Lyn and reached the same conclusion.

5.7. Doppler Tomography

The superoutburst of WZ Sge in 2001 enabled time-resolved spectroscopy during a WZ Sge-type superoutburst. Baba et al. (2002) was able to detect double-peaked emission lines of HeII and the constructed Doppler tomogram showed a spiral pattern. This finding had long been considered as evidence (Kato 2002) of the spiral structure, which is expected by the 2:1 resonance model.

The 2007 superoutburst of V455 And provided another




Fig. 14. Nightly variation of early superhumps in WZ Sge (2001). The data are from Kato et al. (2009a) (same as Ishioka et al. 2002). The numbers in the upper end of upper panels are BJD center -2452100 . The intensity units for the lower panels correspond to 1000 for 8.0 mag.

opportunity to obtain time-resolved high-dispersion spectroscopy with Subaru telescope. The result was quite unexpected with singly peaked emission lines (Nogami et al. 2009). This result could not be easily understood from the 2:1 resonance model. Uemura et al. (2015) reported that obscuration of the inner part of the disk by the disk rim is insufficient to reproduce the observation.

6. Ordinary Superhumps

6.1. Transition from Early to Ordinary Superhumps

In WZ Sge (2001), ordinary superhumps (stage A superhumps, see later sections) smoothly developed from one of two peaks of early superhumps (figure 126 in Kato et al. 2009a). No other object has such extensive coverage and sufficient amplitudes of early superhumps, and it has not yet been confirmed whether such smooth transition is usual for WZ Sge-type dwarf novae.

6.2. Period Variation

The periods of superhumps are known to systematically vary. Kato et al. (2009a) was the first to demonstrate that these period variations have a common pattern in SU UMa-type dwarf novae: initial growing stage (stage A) with a long period and fully developed stage (stage

B) with a systematically varying period and later stage C with a shorter, almost constant period. In WZ Sge-type dwarf novae, stage C is usually not present.

Although stage A superhumps are currently understood to reflect the dynamical precession rate of the disk at the radius of 3:1 resonance (Osaki, Kato 2013b; Kato, Osaki 2013b) and stage B is considered to have a smaller precession rate due to the pressure effect, which produces retrograde precession, in the disk (Osaki, Kato 2013b; Kato, Osaki 2013a), the origin of stage C superhumps and why stage B–C transition suddenly occurs are unsolved problems. During stage B, SU UMa-type dwarf novae show smooth period variations with a more or less constant $P_{\dot{\text{d}}ot} = \dot{P}/P$ (Kato et al. 2009a). Most of systems with short (less than 0.065 d) orbital periods are known to have positive $P_{\dot{\text{d}}ot}$ for stage B (cf. Kato et al. 2009a). Although this variation was originally attributed to an expansion of the disk or an outward propagation of the eccentric wave (Kato et al. 2009a, Howell et al. 1996, Nogami et al. 1998, Baba et al. 2000).⁵ Alternatively, Kato, Osaki (2013a) suggested that the relative strength of the pressure effect

⁵ This idea was originally proposed as a preprint form in 1997 on EG Cnc by Kato et al., and it was introduced in Kato et al. (1998). See Baba et al. (2000) for a detailed description of the background.

Table 2. Amplitudes of Early Superhumps

Object	Orbital period (d)	A1*	A2 [†]	Reference
V455 And	0.05627	0.26	0.22	Kato et al. (2009a)
V466 And	0.05636	0.07	0.038	Kato et al. (2009a)
V500 And	0.05550	–	0.03	Kato et al. (2009a)
V572 And	0.054868	–	0.07	Imada et al. (2006)
EG Cnc	0.05994	–	0.018	Patterson et al. (1998), Matsumoto et al. (1998)
AL Com	0.05667	0.06	0.04	Kato et al. (1996), Patterson et al. (1996)
V1251 Cyg	0.07433	–	0.018	Kato et al. (2009a)
DV Dra	0.05883	–	0.13	Kato et al. (2009a)
PR Her	0.05422	–	0.053	Kato et al. (2013a)
V592 Her	0.05610	–	0.01	Kato et al. (2010)
RZ Leo	0.076038	–	0.05	Kato et al. (2009a)
GW Lib	0.05332	0.00	0.00	Kato et al. (2009a)
SS LMi	0.056637	–	0.15	Shears et al. (2008)
EZ Lyn	0.059005	0.07	0.067	Kato et al. (2012a)
GR Ori	0.058333 [‡]	0.00	0.00	Kato et al. (2014b)
BW Scl	0.054323	0.16	0.10	Kato et al. (2013a)
WZ Sge	0.056670	0.19	0.14	Kato et al. (2009a)
UW Tri	0.05334	–	0.05	Kato et al. (2009a)
CT Tri	0.05281	–	0.03	Kato et al. (2009a)
BC UMa	0.06258	–	0.04	Maehara et al. (2007)
V355 UMa	0.058094	–	0.01	Kato et al. (2012a)
HV Vir	0.057069	0.052	0.044	Kato et al. (2009a)
ASAS SN-13ax	0.056155 [‡]	–	0.00	Kato et al. (2014b)
ASAS SN-14cl	0.05838	–	0.018	Kato et al. (2015)
ASAS SN-14gx	0.05488	–	0.03	Kato et al. (2015)
ASAS SN-14jf	0.05539	–	0.04	Kato et al. (2015)
ASAS SN-15ah	0.05547 [‡]	–	0.00	Kato et al. (2015)
ASAS SN-15bp	0.05563	–	0.014	Kato et al. (2015)
CRTS J090239	0.05652	–	0.35	Kato et al. (2009a)
CRTS J104411	0.05909	–	0.030	Kato et al. (2010)
CRTS J223003	0.05841	–	0.033	Kato et al. (2010)

*Amplitude near the peak brightness.

[†]Mean amplitude.

[‡]Superhump period.

and the dynamical precession by the gravitational field of the secondary may play a key role: lower- q systems have smaller dynamical precession rates and the retrograde precession by the pressure effect becomes relatively more important, making the stronger period variation. However, this explanation was not sufficient to reproduce the long superhumps period at the end of stage B without introducing an expansion of the disk. The physical origin of positive P_{dot} for stage B is still poorly understood.

WZ Sge-type dwarf novae have the same characteristics as in SU UMa-type dwarf novae and extreme WZ Sge-type dwarf novae tend to have smaller P_{dot} . It has been demonstrated that P_{dot} and P_{orb} are correlated with the rebrightening type (starting with figure 36 in Kato et al. 2009a and refined in Kato et al. 2009a–Kato et al. 2014a). In figure 17, we show the updated result up to Kato et al. (2015) (we also used table 3, which shows parameters of the objects in Kato et al. 2015 in the same format as in Kato et al. 2014a). The general tendency is: type-A

outbursts (long rebrightenings) and type-D outbursts (no rebrightening) tend to occur in objects with short P_{orb} . While type-A outbursts usually have small P_{dot} , type-D outbursts can have larger P_{dot} . Type-C outbursts are usually seen in objects with longer P_{orb} and larger P_{dot} , and objects with type-C outbursts are closer to ordinary SU UMa-type dwarf novae. Type-B outbursts usually occur in objects with intermediate P_{orb} and these objects tend to show relatively small P_{dot} . Nakata et al. (2013b) studied two objects with type-B outbursts and found that they occupy a limited region on the $P_{\text{orb}}-P_{\text{dot}}$ diagram. We will discuss this issue later (subsections 7.5, 7.6).

6.3. Delay Time of Superhump Appearance

In figure 18, we show the distribution of the delay time of superhump appearance using the data up to Kato et al. (2015). Although we selected outbursts which were detected sufficiently close to the peak brightness, we should note that actual outbursts may have started slightly ear-

Table 2. Amplitudes of Early Superhumps (continued)

Object	Period (d)	A1*	A2†	Reference
CSS J174033	0.045048	0.033	0.030	T. Ohshima et al. in preparation
MASTER J005740	0.056190	–	0.23	Kato et al. (2014a)
MASTER J085854	0.05556	–	0.00	Kato et al. (2015)
MASTER J094759	0.05588	–	0.006	Kato et al. (2014a)
MASTER J181953	0.05684	–	0.022	Kato et al. (2014a)
MASTER J203749	0.06062	–	0.036	Nakata et al. (2013b)
MASTER J211258	0.05973	–	0.050	Nakata et al. (2013b)
OT J012059	0.057157	–	0.045	Kato et al. (2012a)
OT J030929	0.05615	–	0.018	Kato et al. (2015)
OT J111217	0.05896	–	0.14	Kato et al. (2009a)
OT J112619	0.05423	–	0.04	Kato et al. (2014a)
OT J184228	0.07168	–	0.005	Kato et al. (2013a)
OT J210950	0.05865	–	0.00	Kato et al. (2013a)
OT J213806	0.05450	–	0.04	Kato et al. (2010)
OT J230523	0.05456	–	0.035	Kato et al. (2015)
OT J232727	0.05277	–	0.018	Kato et al. (2014a)
PNV J062703	0.05787	–	0.02	Kato et al. (2014a)
PNV J172929	0.05973	–	0.015	Kato et al. (2015)
SDSS J161027	0.05965	–	0.05	Kato et al. (2010)
SDSS J172325	0.05920‡	–	0.00	Kato et al. (2015)
TCP J153756	0.06101	–	0.038	Kato et al. (2014a)

*Amplitude near the peak brightness.

†Mean amplitude.

‡Superhump period.

Table 3. Parameters of WZ Sge-type superoutbursts in Kato et al. (2015).

Object	Year	P_{SH}	P_{orb}	P_{dot}^*	err*	ϵ	Type†	$N_{reb}‡$	delay§	Max	Min
FI Cet	2014	0.056911	0.05594	9.7	2.1	0.017	–	–	5	14.4	21.6
ASASSN-14cl	2014	0.060008	0.05838	8.5	0.4	0.028	D	0	6	10.7	18.8
ASASSN-14cq	2014	0.057354	0.05660	4.6	0.4	0.013	–	–	8	13.7	21.3:
ASASSN-14cv	2014	0.060413	0.059917	0.9	0.9	0.008	B	8	14	11.2	19.2
ASASSN-14gx	2014	0.056088	0.05488	5.1	0.6	0.022	–	–	9	14.9	21.7:
ASASSN-14jf	2014	0.055949	0.05539	1.1	0.2	0.010	–	–	9	13.3	21.0:
ASASSN-14jq	2014	0.055178	–	4.3	1.2	–	A	–	–]13.7	20.5
ASASSN-14jv	2014	0.055102	0.05442	4.9	0.7	0.013	D	0	9	11.3	19.3
ASASSN-14mc	2014	0.055463	–	1.7	2.1	–	–	–	10	14.3	21.0:
ASASSN-15ah	2015	0.055469	–	6.2	3.2	–	–	–	8	13.6	21.8:
ASASSN-15bp	2015	0.056702	0.05563	4.5	0.3	0.019	B?	–	8	13.6	21.8:
MASTER J085854	2015	0.055560	–	8.1	1.0	–	B	2]4]13.7	18.6:
OT J030929	2014	0.057437	0.05615	6.8	0.5	0.023	D	0	6	11.0	18.9
OT J230523	2014	0.055595	0.05456	8.2	1.3	0.019	–	–	6	12.3	19.8
PNV J171442	2014	0.060092	0.059558	4.4	0.7	0.009	B	5	12	13.5	20.2
PNV J172929	2014	0.060282	0.05973	2.6	1.2	0.009	D	0	11	12.1	21.5

*Unit 10^{-5} .

†A: long-lasting rebrightening; B: multiple rebrightenings; C: single rebrightening; D: no rebrightening.

‡Number of rebrightenings.

§Days before ordinary superhumps appeared.




Fig. 17. $P_{\dot{}}$ versus P_{orb} for WZ Sge-type dwarf novae. Symbols represent the type of outburst: type-A (filled circles), type-B (filled squares), type-C (filled triangles), type-D (open circles) and type-E (filled stars) (see text for details). On the right side, we show mass ratios estimated using equation (6). We can regard this figure as to represent an evolutionary diagram (see discussion in subsection 7.5).

lier. These values should better be considered as lower limits although the difference is likely less than an order of a few days. The delay time of WZ Sge-type objects as a whole has a maximum around 10 d. Type-A outbursts appear to be concentrated around this maximum. Type-D outbursts are more widely distributed, although type-D outbursts may be less favorably observed outbursts (i.e. rebrightening was simply missed due to the faintness of the object and so on), and some of the delay times may have been underestimated more strongly than other types. Type-C outbursts appear to have shorter delay times. Type-B outbursts appear to have a bimodal distribution, below 5 d and more than 12 d. Since these type-B outbursts with short delay times (UZ Boo and EZ Lyn) have been well-examined in order to avoid underestimation, this bimodal distribution appears to be real. The longest delay time (21 d) was seen in OT J111217, which has a very high outburst amplitude (9.4 mag), the second largest in our sample (subsection 3.3). This object appears to be an extreme object in these two respects of statistics.

During this delay time, we can see early superhumps in many systems (section 5). The systematic difference of the delay time between different SU UMA-type dwarf novae with different outburst activity was pointed out by Osaki (1995a), who discussed that this difference may reflect the $1/q^2$ -type dependence of the growth time of the 3:1 resonance (Lubow 1991a; Lubow 1991b). Osaki (1995a) also considered a possibility that the viscous depletion time is longer in some orbital parameters to enable the disk to reach the 3:1 resonance. After the firm identification of early superhumps in WZ Sge, Osaki, Meyer (2003) regarded as the suppression of the 3:1 resonance by the 2:1 resonance Lubow (1991a) as the main cause of the long delay times in WZ Sge-type objects. This interpretation has




Fig. 18. Delay time of superhump appearance. The numbers based on the outburst type is shown. Type-B outbursts appear to show a bimodal distribution.




Fig. 19. Delay time of superhump appearance versus superhump period (as a proxy to orbital period).

been reinforced by the discovery of a faint superoutburst of AL Com in 2015 which did not show early superhump but showed a quick growth of superhumps (Kimura et al. 2015).

Following this interpretation, the delay time reflects the strength of the 2:1 resonance. The statistics of short delay times in type-C outbursts is in line with the interpretation that type-C objects are closer to ordinary SU UMA-type dwarf novae (subsection 6.2). It is interesting that type-B outbursts appear to have two types with different delay times. The objects with shorter delay times appear to have critically reached the 2:1 resonance and this may be in line with the conclusion that objects with type-B outbursts are not (necessarily) good candidates for period bouncers (Nakata et al. 2013b) (see subsection 7.6).

Figure 19 shows the relation between the superhump period (as a proxy to the orbital period) and delay time. This figure also shows that delay times are shorter (i.e. the strength of the 2:1 resonance is weaker) in systems with longer superhump (or orbital) periods.

6.4. Late-Stage Superhumps

Well-observed WZ Sge-type dwarf novae have long persistence of superhumps well after the termination of superoutbursts (e.g. WZ Sge: Patterson et al. 2002, Kato et al. 2008; EG Cnc: Patterson et al. 1998, GW Lib and V455 And: Kato et al. 2008). Probably the best established case is GW Lib in 2007 (Kato et al. 2008 and figure 33 in Kato et al. 2009a), which showed persistent superhumps for at least 930 cycles (50 d) after the termination of the superoutburst. These superhumps showed very constant profile and did not show a phase jump. Patterson et al. (2002) reported the persistence of superhumps in WZ Sge for 90 d (including the outburst part). V355 UMa also showed persistent superhumps for at least 490 cycles (28 d) (Kato et al. 2012a).

These superhumps generally have longer periods than the superhump periods during superoutburst (Kato et al. 2008). Although Kato et al. (2008) interpreted that this increase of the period reflects the expansion of the disk after the outburst, this interpretation was probably incorrect since the pressure effect was not properly treated when comparing superhump periods. As introduced in subsection 6.2, the pressure effect shortens the superhumps period and this effect is strongest in the initial part of stage B (Osaki, Kato 2013b; Kato, Osaki 2013a). Since the identification of stage A superhumps as superhumps reflecting the dynamical precession rate at the 3:1 resonance (Osaki, Kato 2013b; Kato, Osaki 2013b), we could estimate the disk radius in WZ Sge-type dwarf novae after the superoutburst (Kato, Osaki 2013b). The result was $0.37\text{--}0.38A$, where A is the binary separation, in systems without rebrightenings (type-D outbursts) and $0.30\text{--}0.32A$ in systems with long or multiple rebrightenings (type-A or B outbursts) (Kato, Osaki 2013b). There are no measurable samples for type-C outbursts. These experimentally determined disk radius can be used to estimate q for objects (e.g. Kato et al. 2013b).

Transitions from superhumps during the superoutburst plateau to post-superoutburst superhumps are often associated with a disturbance in the $O - C$ diagram (e.g. GW Lib: figure 33 in Kato et al. 2009a; FL Psc = ASAS J002511+1217.2: figure 34 in Kato et al. 2009a). In the case of FL Psc, two hump maxima appeared during the post-superoutburst phase before the rebrightening, and the one peak (0.5 phase different from the superhumps during the superoutburst plateau) smoothly continued as late-stage superhumps. There was also a phase 0.5 jump around the termination of the superoutburst in V355 UMa (Kato et al. 2012a). This phenomenon appears to corresponds to “traditional” late superhumps (superhumps with a ~ 0.5 phase superhumps shift seen during the very late or post-superoutburst stages: e.g. Vogt 1983). In ordinary SU UMa-type dwarf novae, most of originally supposed ~ 0.5 phase shift after the termination of the superoutburst were a result of a combination of incorrect cycle counts and stage C superhumps, which have $\sim 0.5\%$ shorter periods than stage B superhumps (Kato et al. 2009a). Many well-observed SU UMa-type dwarf

novae, including Kepler observations of V585 Lyr (Kato, Osaki 2013a), have been confirmed show continuous $O - C$ diagrams (no phase jump) and the appearance of superhumps with ~ 0.5 phase shift is only limited to high mass-transfer systems [e.g. V344 Lyr (Wood et al. 2011), YZ Cnc (Kato et al. 2014a), V1159 Ori (Patterson et al. 1995), VW Hyi (van der Woerd et al. 1988)]. This observation is in line with the classical interpretation that late superhumps arise from the hot spot on an elliptical disk (Osaki 1985). The presence of a ~ 0.5 phase shift in WZ Sge-type objects, which are considered to be low mass-transfer systems, is a mystery.

The long persistent superhumps in WZ Sge-type object has been considered a result of low mass-transfer rate from the secondary, making the eccentric disk structure to survive longer (Osaki et al. 2001). We should note, however, recent Kepler observations and high-quality ground-based observations have shown that late-stage superhumps persist longer (one or two outburst cycles) after the termination of the superoutburst (e.g. Still et al. 2010; Wood et al. 2011; Osaki, Kato 2013b; Osaki, Kato 2014; Kato et al. 2014b; Kato et al. 2014a) in ordinary SU UMa-type dwarf nova with high mass-transfer rates in contrast to textbook descriptions (e.g. Warner 1995).

6.5. Orbital Variation during Outburst?

Besides the claim of the enhanced orbital signal during rebrightenings in WZ Sge (Patterson et al. 2002) (see subsection 8.1 for the discussion), there was possible transient appearance of the orbital signal in the low-inclination system GW Lib near the end of the superoutburst plateau (Kato et al. 2009a). This phenomenon cannot be easily explained by an enhanced hot spot or by superhump-type modulations.

7. Mass Ratios, Evolutionary Status and Related Topics

7.1. Past Study

CVs evolve as the system lose the total angular momentum and the mass is transferred from the secondary as a result (for recent reviews of CV evolution, see e.g. Kolb, Baraffe 1999; Araujo-Betancor et al. 2005b; Knigge et al. 2011). The orbital period generally decreases during the CV evolution. When the secondary star in the system becomes degenerate, the system reaches the “period minimum” and the orbital period then increases. This is due to two reasons: the thermal time-scale of the secondary exceeds the mass-transfer time-scale and the mass-radius relation is reversed for degenerate dwarfs. The systems evolved beyond the period minimum are usually called “period bouncers”. WZ Sge-type objects have been long considered as candidate period bouncers, and there also have been a discussion whether the secondary is indeed a brown dwarf (cf. Patterson 2011). Currently there are indeed several eclipsing systems having secondary masses in the range of brown dwarfs (SDSS J103533.02+055158.3: Littlefair et al. 2006, Savoury et al. 2011, OV Boo = SDSS J150722.33+523039.8: Littlefair et al. 2006, SDSS

J143317.78+101123.3: Savoury et al. 2011, PHL 1445 = PB 9151: McAllister et al. 2015), although none of them have been confirmed to be a WZ Sge-type object by undergoing an outburst. Patterson et al. (1998), Patterson (1998), Patterson (2011) used fractional superhump excesses to estimate the mass ratios, and argued that some of WZ Sge-type dwarf novae are good candidates for period bouncers.

EG Cnc, which displayed six rebrightenings, was particularly notable in the small estimated $q = 0.035$ (Patterson et al. 1998). WZ Sge itself was also claimed to have a small $q = 0.045$ (Patterson et al. 2002) by the same method. Patterson et al. (2002) also suggested that the secondary was very faint in the infrared, supporting the brown-dwarf hypothesis. Steeghs et al. (2001), however, derived $q = 0.040\text{--}0.075$ by Doppler tomography. Harrison et al. (2013) also suspected a high fraction of infrared emission from the secondary. The situation remained unclear whether traditionally known WZ Sge-type objects are period bouncers or not.

7.2. Determination of Mass Ratios using Stage-A Superhumps

Following the identification of stage A superhumps reflecting the dynamical precession at the radius of the 3:1 resonance (subsection 6.2), we have now become able to measure mass ratios of WZ Sge-type objects (Kato, Osaki 2013b). We briefly review the outline of the method here. The precession rate of the disk, $\omega_{\text{dyn}}/\omega_{\text{orb}}$, is equivalent to the fractional superhump excess (in frequency) $\epsilon^* \equiv 1 - P_{\text{orb}}/P_{\text{SH}}$ and it is related to the conventional fractional superhump excess (in period) $\epsilon \equiv P_{\text{SH}}/P_{\text{orb}} - 1$ by a relation $\epsilon^* = \epsilon/(1 + \epsilon)$. The dynamical precession rate ω_{dyn} is

$$\omega_{\text{dyn}}/\omega_{\text{orb}} = Q(q)R(r), \quad (1)$$

where ω_{orb} and r are the angular orbital frequency and the dimensionless radius measured in units of the binary separation A . The dependence on q and r are (cf. Hirose, Osaki 1990)

$$Q(q) = \frac{1}{2} \frac{q}{\sqrt{1+q}}, \quad (2)$$

and

$$R(r) = \frac{1}{2} \frac{1}{\sqrt{r}} b_{3/2}^{(1)}(r), \quad (3)$$

where $\frac{1}{2}b_{s/2}^{(j)}$ is the Laplace coefficient

$$\frac{1}{2}b_{s/2}^{(j)}(r) = \frac{1}{2\pi} \int_0^{2\pi} \frac{\cos(j\phi)d\phi}{(1+r^2-2r\cos\phi)^{s/2}}. \quad (4)$$

Considering that the superhump wave is confined to the 3:1 resonance region during stage A (hence the pressure effect can be neglected and the precession frequency reflects the pure dynamical one), we can substitute r by the radius of the 3:1 resonance.

$$r_{3:1} = 3^{(-2/3)}(1+q)^{-1/3}, \quad (5)$$

Then $Q(q)R(r_{3:1})$ becomes a function of q and we can directly estimate q from ϵ^* of stage A superhumps.

This method is particularly useful for WZ Sge-type objects, since they usually show early superhumps, which have periods almost identical with the orbital periods, and stage A superhumps develop immediately following the stage of early superhumps. After a typical waiting time of ~ 10 d (subsection 6.3), we can relatively easily detect stage A superhumps and determine mass ratios. In papers Kato, Osaki (2013b), Nakata et al. (2013b), Kato et al. (2014b), Kato et al. (2014a), Kato et al. (2015), a sizable number of WZ Sge-type objects have been determined for q using this method. The most up-to-date evolutionary diagram is shown in Kato et al. (2015).

Kato, Osaki (2013b) also demonstrated that traditional methods (such as Patterson 1998; Patterson 2011) for determining q using (stage B) superhumps during the superoutburst plateau give systematically small q values for small- q systems because the pressure effect decreases the precession rate of this eccentric disk, and this relative importance of the effect is larger for systems with smaller precession rates (i.e. WZ Sge-type objects). This explains why Patterson (2011) listed so many candidates for period bouncers using the fractional superhump excess. See Kato, Osaki (2013b) for more detailed discussion.

7.3. Current Understanding

This new method has clarified mass ratios of many WZ Sge-type objects and clarified the evolutionary path to an unprecedented detail. According to the estimates in Kato, Osaki (2013b), many WZ Sge-type dwarf novae (WZ Sge itself, too) have mass ratios near the borderline between lower main-sequence and brown dwarf secondaries. The most recent work (Kato et al. 2015) indicates a high concentration of WZ Sge-type object around orbital periods of 0.054–0.056 d and mass ratios 0.06–0.08. The spread of mass ratios in this region and the absence of objects in shorter periods suggests that these objects are indeed near the period minimum, and WZ Sge-type objects are indeed located near the period minimum. Now it is no wonder some objects (such as WZ Sge, Harrison et al. 2013) have some evidence of infrared emission from the secondary while other objects have either undetectable secondaries or brown dwarf secondaries have been identified by eclipse studies in WZ Sge-type candidates (subsection 7.1).

It looks like that “prototypical” WZ Sge-type objects such as WZ Sge have intermediate mass ratios among the WZ Sge-type objects. More unusual objects (such as higher outburst amplitudes) seem to have lower mass ratios.

7.4. Distribution of Mass Ratios

Figure 20 shows the distribution of ratios in WZ Sge-type objects and non-WZ Sge-type objects using stage A superhump method. The used data up to Kato et al. (2015) are summarized in table 4. There is a sharp peak between $q = 0.07$ and $q = 0.08$. The mean and standard deviation of this distribution are 0.078 and 0.017, respectively. Although it is difficult to define the upper limit

Table 4. Mass Ratios of WZ Sge-Type Objects from Stage A Superhumps

Object	Oribital period (d)	ϵ^*	error	q	error	$P_{\text{dot}} \times 10^5$	error
from Kato et al. (2009a)							
V455 And	0.05631	0.0296	0.0014	0.080	0.004	4.7	1.2
V466 And	0.05636	0.0308	0.0013	0.083	0.004	5.7	0.7
HO Cet	0.05490	0.0328	0.0012	0.090	0.004	4.9	0.5
GW Lib	0.05332	0.0258	0.0014	0.069	0.003	4.0	0.1
WZ Sge	0.05669	0.0290	0.0010	0.078	0.003	2.0	0.4
HV Vir	0.05707	0.0268	0.0003	0.072	0.001	6.8	0.4
ASAS J102522	0.06136	0.0423	0.0018	0.120	0.005	10.9	0.6
from Kato et al. (2010)							
V592 Her	0.05610	0.0206	0.0049	0.054	0.014	7.4	0.6
SDSS J161027	0.05687	0.033	0.0015	0.090	0.005	6.8	1.4
CRTS J104411	0.05909	0.0288	0.0048	0.077	0.001	–	–
from Kato et al. (2012a)							
EZ Lyn	0.05901	0.0290	0.0011	0.078	0.003	3.5	0.9
V355 UMa	0.05729	0.0247	0.0005	0.066	0.001	5.4	0.2
from Kato et al. (2013a)							
BW Scl	0.05432	0.0251	0.0021	0.067	0.006	4.4	0.3
OT J184228	0.07168	0.0163	0.0007	0.042	0.003	-0.9	1.5
OT J210950	0.05865	0.0365	0.0010	0.101	0.003	8.5	0.6
from Kato et al. (2014b)							
MASTER J094759	0.0559	0.0225	0.0029	0.059	0.008	3.0	1.1
MASTER J181953	0.05684	0.0259	0.0003	0.069	0.001	2.6	1.1
MASTER J211258	0.05973	0.0300	0.0005	0.081	0.002	0.8	1.0
OT J112619	0.05423	0.0317	0.0006	0.086	0.002	3.6	0.4
OT J203749	0.06051	0.0351	0.00011	0.097	0.008	2.9	1.0
OT J232727	0.05277	0.0303	0.0005	0.082	0.002	4.0	1.1
SSS J122221	0.075879	0.0172	0.0001	0.044	0.001	-1.1	0.7
from Kato et al. (2014a)							
MASTER J005740	0.056190	0.0280	0.0050	0.076	0.016	4.0	1.0
PNV J191501	0.05706	0.0344	0.0012	0.095	0.004	5.2	0.2
TCP J233822	0.057255	0.0231	0.0014	0.061	0.004	2.7	1.1

Table 4. Mass Ratios of WZ Sge-Type Objects from Stage A Superhumps (continued)

Object	Oribital period (d)	ϵ^*	error	q	error	$P_{\text{dot}} \times 10^5$	error
from Kato et al. (2015)							
ASASSN-14cv	0.059917	0.0286	0.0003	0.077	0.001	0.9	0.9
ASASSN-14jf	0.05539	0.0260	0.0020	0.070	0.005	1.1	0.2
ASASSN-14jv	0.05442	0.0278	0.0009	0.074	0.003	4.9	0.7
ASASSN-15bp	0.05563	0.0293	0.0006	0.079	0.002	4.5	0.3
OT J030929	0.05615	0.0291	0.0003	0.078	0.001	6.8	0.5
OT J213806	0.054523	0.041	0.0004	0.120	0.020	7.2	0.4
OT J230523	0.05456	0.0366	0.0007	0.102	0.002	8.2	1.3
PNV J171442	0.059558	0.0284	0.0003	0.076	0.001	4.4	0.7
PNV J172929	0.05973	0.0273	0.0005	0.073	0.002	2.6	1.2




Fig. 20. Distribution of mass ratios in WZ Sge-type objects and non-WZ Sge-type objects using stage A superhump method.

of mass ratios for WZ Sge-type objects, we can choose 0.09 which is close to the one sigma above the mean of WZ Sge-type objects and close to the lower end of non-WZ Sge-type objects. This value may be considered as an empirical limit of mass ratios which enable the 2:1 resonance in outburst. Some objects have apparently higher (above 0.10) mass ratios but still show early superhumps. The limit is probably not a rigid border but probably depends on the strength of the outburst or other factors. It would be interesting to compare this result with figure 2 in Osaki, Meyer (2002). The present limit corresponds to $\log q = -1.0$, which is close to the limit suggested in the dashed line of figure 2 in Osaki, Meyer (2002).

7.5. Period Variation and Mass Ratio

Although the mechanism is not yet clear, superhump periods systematically vary during stage B (subsection 6.2). The values of P_{dot} are strongly related to the orbital periods (cf. Kato et al. 2009a). Since P_{dot} and rebrightening type are empirically known to be strongly related (subsection 6.2), we here examined whether P_{dot} can be used as a measure of q . Figure 21 shows the relation between P_{dot} and q for WZ Sge-type objects using stage A superhump method. Although some data points have large errors, the overall appearance suggests that P_{dot} is almost linearly related to q at least for WZ Sge-type objects. This relation strengthens our impression (in our series papers up to Kato et al. 2015) that WZ Sge-type objects having properties similar to those of SU UMa-type dwarf novae have larger P_{dot} and extreme WZ Sge-type objects have smaller P_{dot} . We have derived

$$q = 0.0043(9)P_{\text{dot}} \times 10^5 + 0.060(5). \quad (6)$$

We should note, however, this relation does not necessarily holds in longer period systems (ordinary SU UMa-type dwarf novae) as suggested by the presence of a number of long-period SU UMa-type objects with unusual values of P_{dot} (e.g. Kato et al. 2013a).




Fig. 21. Relation between period variations and mass ratios for WZ Sge-type objects using stage A superhump method.

Assuming this linear relation holds in all WZ Sge-type objects, figure 17 can be directly read as a diagram between P_{orb} , q and the outburst type. The y -axis of figure 17 then corresponds to the q range of 0.047–0.11.

7.6. Rebrightening Type and Evolution

As discussed in the previous subsection, we suggest that the $P_{\text{orb}}-P_{\text{dot}}$ diagram can be regarded as an evolutionary diagram. From figure 17, the objects in the upper branch (non-degenerated, low-mass main sequence secondary, close to ordinary SU UMa-type dwarf novae) tend to show type-C outbursts. The system closer to the period minimum but still on the upper branch, type-D outbursts are most frequently met. Around the period minimum, type-A outbursts are more dominant. Type-B outbursts are more widely spread, but at least some of them are good candidates for period bouncers (P_{dot} close to zero or even negative). It appears that objects with type-B outbursts have two populations as already discussed in subsection 6.3. This result is consistent with C. Nakata et al. in preparation, who apparently found a population of objects with type-B outbursts different from Nakata et al. (2013b).

Thanks to the new estimate method of q using stage A superhumps, we can now recognize the outburst type as a kind of evolutionary sequence (type C \rightarrow D \rightarrow A \rightarrow B \rightarrow E, with some outliers for type-B objects). Theoretical interpretation of the relation between q and P_{dot} and of these different types of rebrightenings are eagerly sought.

7.7. Duration of Stage A Superhump Phase

We placed this subsection here since it is most related to the mass-ratio issues. As discussed in subsection 6.3, the growth time of the 3:1 resonance is a dependence of $1/q^2$ (Lubow 1991a; Lubow 1991b), and it was originally suggested for explaining the delay of appearance of superhumps, although Osaki, Meyer (2003) later did not adopt this interpretation for the delay of appearance of superhumps. We can now measure the growth time

of stage A superhumps. We used the same in table 4 (excluding HO Cet, which was not sufficiently sampled). The durations of stage A were estimated from the lower limit of E_1 (start of stage B in cycles) in summary tables such as table 2 in Kato et al. (2009a). Two objects have been added from C. Nakata et al. in preparation (cf. subsection 7.8): OT J230425 with $P_{\text{SH}}=0.06628(6)$ d (stage B), $P_{\text{dot}}=-3.9(2.4)$, duration of stage A = 123 cycles and OT J075418 with $P_{\text{SH}}=0.07076(1)$ d (stage B), $P_{\text{dot}}=-2.4(0.5)$, duration of stage A = 190 cycles. We should note these durations are lower limits rather than firm estimates, since there are usually gaps in observation lasting less than 1 d and it is usually difficult to detect low-amplitude stage A superhumps in the beginning. The estimates, however, are not expected to be shorter by more than 20 cycles (corresponding to ~ 1 d) than the real values. We did not plot these uncertainties in the figures.

The relation period variations and duration of stage A phase for WZ Sge-type objects is shown in figure 22. The upper panel directly shows the comparison between P_{dot} and duration of stage A phase. Objects with lower P_{dot} have longer stage A phases. Considering that P_{dot} is a good measure of q (subsection 7.5), this relation agrees to what is expected. In the lower panel, estimated q values using equation (6) are used. The slope in the log-log diagram is expected to be -2 if the growth time has a $1/q^2$ dependence. The result appears to be consistent with this expectation. We consider that the duration of stage A phase can be a useful probe for estimating the evolutionary phase, even if ϵ^* or P_{dot} is not directly determined.

7.8. Period Bouncers

Period bouncers are CVs past the period minimum. The presence of the period minimum was proposed early (Paczynski, Sienkiewicz 1981). Rappaport et al. (1982) and Paczynski, Sienkiewicz (1983) are early model calculations of the CV evolution. There have been many refined model calculations (Kolb 1993; Kolb, Baraffe 1999; Howell et al. 2001). Although we do not intend to go deep into the problems of CV evolution, it has widely been recognized that evolutionary time of CVs is significantly shorter than the Hubble time, and most of CVs must have already passed the period minimum. This implies there must be many period bouncers.

Observational evidence for this picture had been scarce until SDSS discovered many faint CVs (Gänsicke et al. 2009). Gänsicke et al. (2009) was the first to demonstrate the presence of the “period spike” around the period minimum. However, period bouncers still remained elusive objects. Although eclipse observations have revealed a number of objects which contain secondaries whose masses are comparable to brown dwarfs (see subsection 7.1 for such objects), the majority of the theoretically expected period bouncers remained almost a missing population. Although Patterson (2011) listed candidates based on several criteria, most of the listed objects did not meet sufficiently many criteria. Patterson (2011) listed estimates of q values for some objects using the traditional method for converting fractional superhump excesses to q . As already




Fig. 22. Relation between period variations and duration of stage A phase for WZ Sge-type objects. (Upper) Relation between P_{dot} and duration of stage A phase. Objects with lower P_{dot} have longer stage A phases. (Lower) Relation between q estimated from equation (6) and duration of stage A phase (in log scale).

introduced in subsection 7.2, this traditional method gives systematically small q values for low- q objects like WZ Sge-type objects (Kato, Osaki 2013b), and some of the listed objects in Patterson (2011) may not be good candidates for period bouncers.

Kato et al. (2013b) found that SSS J122221 showed two successive superoutbursts (currently classified as type-E outburst) and that superhumps were developed during the second superoutburst. Stage A superhumps and post-superoutburst superhumps were detected and Kato et al. (2013b) obtained a stringent limit of $q < 0.05$ based on the stage A superhumps and dynamical precession model (subsection 7.2). Combined with the long superhump period (~ 0.0765 d), this object, together with OT J184228 showing type-E outburst, was proposed to be the best candidate for period bouncers.

Following this work, Nakata et al. (2014) identified two objects (OT J075418, OT J230425) as additional good candidates. These objects, together with SSS J122221 and OT J184228, share common properties: (1) extremely long-lasting phase (100–200 cycles) of stage A (subsection 7.7) and (2) very slow fading rates (less than 0.05 mag d^{-1}) (subsection 4.5) and (3) long superhump periods (longer than 0.065 d). We consider they are currently

known best criteria of period bouncers if they undergo superoutbursts. Nakata et al. (2014) also made a statistical consideration assuming that mass-transfer rates (from the secondary) for these systems are 10–100 times lower than ordinary SU UMa-type objects with similar superhump periods, based on the standard evolutionary model assuming the gravitational wave radiation as the main source of angular momentum loss, reached a conclusion that the recently discovered fraction of these candidate period bouncers among SU UMa-type dwarf novae can account for the theoretically expected population of period bouncers. Characterization of these objects by detailed observations is now desired to unveil the nature of period bouncers. A further search for objects (some objects are already proposed in this paper, see section 10) using the above criteria will surely enrich our knowledge in period bouncers and CV evolution.

7.9. Long-Period Objects

There are objects other than (candidate) period bouncers with long orbital periods (approximately longer than 0.07 d, see subsection 3.4) which show early superhumps and classified as WZ Sge-type in this paper. If the secondary is a normal lower main-sequence star, the q values for such systems should fall far outside the upper limit ($q=0.09$ according to subsection 7.4) of the 2:1 resonance. There are two possibilities in such systems: (1) either the white dwarf and or the secondary is anomalous and the true q is smaller, or (2) higher- q systems enable the 2:1 resonance in certain conditions. We consider (2) as an interesting possibility, since these objects have longer recurrence times compared to ordinary SU UMa-type dwarf novae. The outburst amplitude is also large (8.0 mag in V1251 Cyg). We consider that in such systems the mass is more accumulated than in ordinary SU UMa-type dwarf novae and the disk can reach the 2:1 resonance when the outburst is violent enough.

7.10. Absolute Magnitudes in Quiescence

Warner (1987) showed that the absolute magnitudes of outbursting dwarf novae are almost constant following a weak linear function of the orbital period. Patterson (2011) refined the relation using greatly improved statistics and showed that the absolute magnitudes of outbursting dwarf novae are a good “standard candle”. The relation has a theoretical foundation (Osaki 1996; Cannizzo 1998; Smak 2000) assuming that the disk-instability model is responsible for dwarf nova outbursts (section 8).

As we will see in subsection 8.2, WZ Sge-type superoutbursts are different from ordinary SU UMa-type superoutbursts in that there is a phase of viscous decay before ordinary superhumps appear. During this phase, the disk mass is much larger than ordinary SU UMa-type superoutbursts and the maximum brightness is not expected to be a standard candle. We instead used the magnitude when ordinary superhumps appears. As we will see in subsection 8.2, the disk is expected to have a size close to the radius of 3:1 resonance and the condition is analogous to ordinary SU UMa-type superoutbursts. In table 5, we

collected magnitudes of WZ Sge-type objects when ordinary superhumps appear [the data source is observations in Kato et al. (2009a)–Kato et al. (2015)]. The measurement of these magnitudes is usually very easy and has a typical error of ± 0.1 mag. Only the objects with certain quiescent magnitudes (such as SDSS magnitudes) are selected in the table. The order of the table is the same as in table 6. We have added candidate period bouncers at the bottom for comparison purposes even though some of these objects do not have certain quiescent magnitudes.

Since there is no estimate for the absolute magnitude when ordinary superhumps appear in the literature, we describe them as differences from quiescent magnitudes. In figure 23, we show the result together with SU UMa-type objects other than WZ Sge-type objects (we call ordinary SU UMa-type objects in this subsection). Since this figure is only for a comparison purpose, we omit the table and source information for each SU UMa-type object to avoid too much complexity and simply list the objects as the form of a footnote.⁶ In this diagram, ordinary SU UMa-type objects are widespread, while WZ Sge-type objects have surprisingly similar outburst “amplitudes” when superhumps start to appear. The mean and standard deviation of these values (excluding candidate period bouncers) are 6.4 and 0.7 mag, respectively. The values for the SU UMa-type objects are 4.3 and 1.1 mag, respectively. Considering the variability in quiescence and the effect of inclination (we neglected the both effects), it would not be an exaggeration to say that all WZ Sge-type objects essentially have the same outburst “amplitudes” when superhumps start to appear. If the absolute magnitude at the appearance of ordinary superhumps is constant, this implies that the quiescent absolute magnitudes in WZ Sge-type objects are almost the same, which gives somewhat different impression from figure 4 in Patterson (2011).

Another surprise is that these “amplitudes” are only marginally larger (only by 0.8 mag) in period bouncers (mean 7.2 mag and standard deviation 0.6 mag) compared to WZ Sge-type objects. If it is indeed the case, absolute magnitudes in quiescence may not be a useful tool for selecting period bouncers.

We consider the zero point of this figure by two methods. The first one is according to the well-known relation in Warner (1987). The absolute magnitude of the outburst is

$$M_V(\text{max}) = 5.64 - 0.259P_{\text{orb}}(\text{hr}). \quad (7)$$

For P_{orb} of most WZ Sge-type objects, this relation gives $M_V(\text{max})=5.3$ [the dependence on P_{orb} is so small that even if we assume a period of 2.0 hr, $M_V(\text{max})=5.1$].⁷

⁶ KX Aql, VY Aqr, EG Aqr, TT Boo, V342 Cam, V391 Cam, OY Car, GX Cas, HT Cas, WX Cet, Z Cha, PU CMa, YZ Cnc, GO Com, TV Crv, V503 Cyg, KV Dra, MN Dra, AQ Eri, AW Gem, V844 Her, CT Hya, MM Hya, VW Hyi, WX Hyi, RZ LMi, SX LMi, BR Lup, V453 Nor, DT Oct, V1032 Oph, V1159 Ori, V368 Peg, QY Per, AR Pic, TY PsA, V893 Sco, NY Ser, V493 Ser, RZ Sge, SU UMa, SW UMa, SW UMa, BZ UMa, CY UMa, DI UMa, DV UMa, ER UMa, IY UMa, KS UMa, HS Vir, QZ Vir.

⁷ Warner (1987) did not make special distinction between the out-




Fig. 23. Amplitude of outburst at the time of appearance of ordinary superhumps. The filled circles, open circles and filled stars represent WZ Sge-type, SU UMa-type (other than WZ Sge-type) and period bouncers, respectively. The data for WZ Sge-type objects and period bouncers are from table 5. The data for SU UMa-type objects are from various sources, including VSNET data, AAVSO data, CRTS data and SDSS data and periods from Kato et al. (2009a)–Kato et al. (2015). The orbital periods for the objects only with superhump periods were estimated the relation between the orbital and superhump periods (equation 6 in Kato et al. 2012a).

According to our assumption, the condition when ordinary superhumps appear is the same both in ordinary SU UMa-type objects and WZ Sge-type objects. If it is indeed the case, we can adopt the zero point of $M_V=5.3$ for this figure.

The second one is to use the WZ Sge-type objects with known parallaxes. The distance of $43.5(\pm 0.3)$ pc for WZ Sge (Harrison et al. 2004; Thorstensen 2003), $104(+30, -20)$ pc for GW Lib (Thorstensen 2003), and 74 pc for V455 And (listed in Patterson 2011), give zero points of 6.7, $5.1(\pm 0.6)$ and 6.6, respectively. Since WZ Sge and V455 And are high-inclination systems, and GW Lib is nearly a pole-on system (Thorstensen et al. 2002), these values need to be corrected. By using the corrections in table 1 in Patterson (2011), these values become 5.5, $6.0(\pm 0.6)$ and 5.9, respectively. Considering the uncertainties, the zero point of the figure will be in the range of $M_V=5.3$ and $M_V=5.8$. This value is close to the absolute magnitude of the plateau phase of $M_V=5.5(\pm 0.2)$ in Patterson (2011).

We consider the magnitude when ordinary superhumps start to appear is an excellent index (for estimating the distance or the absolute quiescent magnitude) since it is easily defined observationally and can be easily measured to an accuracy of 0.1 mag.

burst types. The values of $m_V(\max)$ for SU UMa-type dwarf novae are in agreement with our magnitudes when superhumps start to appear within 0.5 mag.

Table 5. Brightness when Superhumps Appear.

Object	Year	Mag1*	Mag2†	P_{orb}^\ddagger
WZ Sge	2011	9.9	15.3	0.05669
AL Com	1995	13.6	19.8	0.05667
AL Com	2001	13.5	19.8	0.05667
AL Com	2013	13.6	19.8	0.05667
EG Cnc	1996	12.5	18.0	0.05997
HV Vir	1992	13.3	19.2	0.05707
HV Vir	2002	13.2	19.2	0.05707
HV Vir	2008	13.4	19.2	0.05707
RZ Leo	2000	12.7	18.7	0.07603
RZ Leo	2006	12.6	18.7	0.07603
QZ Lib	2004	12.2	18.8	0.06460s
UZ Boo	2003	12.8	19.7	0.0620s
UZ Boo	2013	12.7	19.7	0.0620s
V592 Her	1998	14.7	21.4	0.0561
V592 Her	2010	14.6	21.4	0.0561
ASAS J102522	2006	12.5	19.3	0.06136
EZ Lyn	2010	12.6	17.8	0.05901
GW Lib	2007	10.2	17.2	0.05332
V455 And	2007	10.9	16.1	0.05631
OT J111217	2007	14.4	20.9	0.05847
SDSS J161027	2009	14.6	20.1	0.05687
CRTS J104411	2010	13.7	19.3	0.05909
OT J012059	2010	14.2	20.1	0.05716
V355 UMa	2011	10.8	17.7	0.05729
OT J210950	2011	12.2	18.7	0.05865
SV Ari	2011	15.0	22.1	0.05552s
BW Scl	2011	10.7	16.5	0.05432
PR Her	2011	13.7	21.0	0.05422
MASTER J211258	2012	15.2	21.3	0.05973
OT J232727	2012	14.9	21.8	0.05277
MASTER J081110	2012	15.0	22.1	0.05814s
OT J112619	2013	15.8	21.8	0.05423

*Brightness when ordinary superhumps appear.

†Quiescent brightness.

‡Orbital or superhump (s) period.

8. Outburst Mechanism

8.1. History

The high-amplitude of the outburst (typically 8 mag), extremely long cycle length (22–33 yr in WZ Sge) and the long duration of the outburst were main features which puzzled researchers. The WZ Sge-type objects became recognized just in pace with the development of the disk instability (DI) model: Osaki 1974; Hōshi 1979; Meyer, Meyer-Hofmeister 1981) and there were heated debates between the mass-transfer burst (MTB) model (originally Bath 1973). In recent years, there have been a wide consensus that the DI model generally accounts for the outburst phenomenon of dwarf novae (cf. Warner 1995; Hellier 2001a). After the debates over the cause of dwarf nova-outburst settled, the next target of debates became superoutbursts of SU UMa-type dwarf novae. Osaki (1985) was the first to propose irradiation-

Table 5. Brightness when Superhumps Appear (continued).

Object	Year	Mag1*	Mag2†	$P_{\text{orb}}‡$
GR Ori	2013	14.7	22.4	0.05833s
MASTER J165236	2013	15.7	22.1	0.08473
MASTER J181953	2013	15.0	21.6	0.05684
PNV J191501	2013	11.7	18.5	0.05706
ASASSN-13ax	2013	14.8	21.2	0.05616s
ASASSN-13ck	2013	14.8	20.8	0.05535
TCP J233822	2013	14.9	21.5	0.05726
MASTER J005740	2013	16.7	20.9	0.05619
ASASSN-14ac	2014	15.3	21.6	0.05855s
PNV J172929	2014	14.3	21.5	0.05973
ASASSN-14cl	2014	11.9	18.8	0.05838
ASASSN-14cv	2014	12.9	19.2	0.05992
FI Cet	2014	15.5	21.6	0.05594
OT J230523	2014	13.4	19.8	0.05456
OT J030929	2014	12.2	18.9	0.05615
ASASSN-14jv	2014	12.5	19.3	0.05442
ASASSN-15bp	2014	13.7	20.5	0.05563
Long-period and borderline systems				
V1251 Cyg	2008	13.5	20.5	0.07433
BC UMa	2003	12.6	18.5	0.06261
Candidate period bouncers				
OT J230425	2010	13.7	21.1	0.06628s
OT J184228	2011	13.6	20.6	0.07168
SSS J122221	2013	12.3	18.8	0.07649s
OT J075418	2013	14.9	22.8	0.07076s
OT J060009	2014	12.9	20.2	0.06331s

*Brightness when ordinary superhumps appear.

†Quiescent brightness.

‡Orbital or superhump (s) period.

induced mass-transfer variation as the cause of super-outbursts. Although this author retracted this idea by proposing the thermal-tidal instability (TTI) model for SU UMa-type dwarf novae (Osaki 1989), the modified MTB-type idea has been studied by many researchers (most notably, Smak 1991, Smak 2004 are representative recent papers).

Most recently, Osaki, Kato (2013a) used Kepler data for V1504 Cyg and succeeded in demonstrating the disk radius variation using negative superhumps. The variation of the disk radius exactly confirmed the prediction by the TTI model, and this observation became the best proof for the TTI model for ordinary SU UMa-type dwarf novae.⁸

The situation for WZ Sge-type outbursts is less clear. Patterson et al. (1981) is the first to identify the highly enhanced orbital variation (currently identified as early superhumps) during the initial nights of the 1978–1979 outburst of WZ Sge. Based on the amplitudes of orbital humps, Patterson et al. (1981) estimated an enhancement of the mass-transfer rate during the outburst by a factor of 60–1000. This observation was supportive of the

MTB model for the WZ Sge-type outburst. During the 2001 superoutburst of WZ Sge, Ishioka et al. (2002) discovered that early superhumps developed while the object was still brightening (see also subsection 5.2), which excluded the possibility that *the outburst is initiated by a sudden mass-transfer burst*. After lucid explanation of early superhumps by the 2:1 resonance by Osaki, Meyer (2002), Patterson et al. (2002) considered that the 2:1 resonance is the most promising explanation for early superhumps. Patterson et al. (2002), however, claimed the detection of enhanced orbital humps in the phase-averaged light curves in the later stage, and claimed that they are the best evidence for enhanced mass-transfer in WZ Sge-type outburst. Osaki, Meyer (2003) discussed this issue and they concluded that the enhanced orbital humps are the aspect effect in a high-inclination system by modeling the orbital light curves based on dissipation pattern of the superhump light source. According to Osaki, Meyer (2003), what looked like the enhanced hot spot by eclipse observations by Patterson et al. (2002) is actually the superhump light source.

8.2. Main Outburst

Both DI(TTI)-type model and DI model modified by the MTB-type effect have been proposed to explain outbursts of WZ Sge-type objects. The most serious consequence for the DI-type model is that it requires extremely low quiescent viscosity is needed ($\alpha_C < 0.00005$: Smak 1993; $\alpha_C < 0.003$: Osaki 1995a) to explain the extreme interval (33 yr) of outbursts in WZ Sge by avoiding inside-out outburst caused by viscous diffusion during quiescence. There are two approaches to tackle with this problem: (1) the quiescent viscosity is indeed extremely low in WZ Sge-type objects, and (2) quiescent viscosity is similar to that of ordinary SU UMa-type dwarf novae, but outbursts are suppressed. The approach (1) is by Osaki (1995a), while a representative approach (2) is by Lasota et al. (1995), who considered the truncation of the inner disk either by magnetic fields (Livio, Pringle 1992) or by the coronal syphon flow (Meyer, Meyer-Hofmeister 1994) and avoided thermal instability to occur. The outburst should necessarily be initiated by MTB-type enhanced mass-transfer in the latter model.

Warner et al. (1996) proposed a model in that the inner disk is truncated by the magnetism of the white dwarf and the mass-transfer rate is just above the critical mass-transfer rate. This model did not require an enhanced mass-transfer to produce an outburst.

Although Warner et al. (1996) succeeded in reproducing the long recurrence time, such an approach with a standard α_C have their own problem, that is, the resultant duration of the outburst should be shorter than reality [i.e. the duration should be similar to ordinary SU UMa-type dwarf novae; in the case of Warner et al. (1996), the duration was only 6 d] since the stored mass is roughly inversely proportional to α_C [Osaki (1996); Osaki (1998), who referred to such a disk as a “leaky bucket” in comparison to a low- α_C as a “big bucket”]. Meyer-Hofmeister et al. (1998) also criticized the model by Warner et al.

⁸ The criticisms by Smak (2013) have been confuted by Osaki, Kato (2014).

(1996) based on the short duration of the computed outburst and on the absence of short outbursts preceding superoutburst in actual observation.

In the meantime, the origin of the quiescent viscosity became more apparent, and α_C can become very small in cold accretion disks (such as in the disks with very low mass-transfer rate as in WZ Sge-type objects) since the magnetic fields decay due to finite conductivity in the cold disk (Gammie, Menou 1998). This interpretation was proposed by Osaki et al. (2001), who considered a model to reproduce repetitive rebrightenings in EG Cnc.

Although there have recently been less arguments against the low- α_C model after the development of our knowledge about the origin of viscosity in accretion disks, attempts with standard α_C in line with the idea by Warner et al. (1996) have been sought [e.g. “magnetic propeller” model in Matthews et al. (2007)]. For a detailed comparison of different models, see Meyer-Hofmeister et al. (1998).

In the TTI model, the WZ Sge-type outburst is explained in the following way (Osaki 1995a; Osaki, Meyer 2003). Since α_C is sufficiently low [the model by Osaki (1995a) considered no viscosity in quiescence, and Meyer-Hofmeister et al. (1998) considered the consequences of a finite viscosity], the transferred matter does not spread by diffusion and stored in the outer part of the disk. This condition avoids the matter to spread to the inner disk and cause an inside-out outburst, which is not supported by observation. The accumulation time is long enough, and the stored angular momentum is sufficient to expand the disk beyond the 3:1 resonance once outburst occurs and all outbursts become superoutbursts (cf. Ichikawa, Osaki 1994; Osaki 1996). Since the stored mass in the disk is large, the outburst starts with a long viscous decay (subsection 4.1), and when the 3:1 resonance starts working, superhumps develop and the outburst follows the evolution of a superoutburst of ordinary SU UMa-type dwarf novae. The main difference in this model from ordinary SU UMa-type dwarf novae is the absence of normal outbursts and the presence of a long viscous decay at the start of the superoutburst. At the time of Osaki (1995a), early superhumps were not known and the 2:1 resonance was not considered. Later this viscous decay phase is found to be governed by the 2:1 resonance and this resonance suppresses the growth of the 3:1 resonance (subsection 6.3).

8.3. Mechanism of Repeated Rebrightenings

The repeated rebrightenings of EG Cnc (1996–1997) attracted many researchers. Osaki et al. (1997) proposed a working model in which the rebrightenings could be reproduced if α_C remained higher than in quiescence some time after the end of the main superoutburst. Osaki et al. (1997) originally considered the possible source of this high α_C as a result of remaining turbulence in the disk which remains eccentric for a long time after the end of the superoutburst. This model was refined by considering the magneto-hydrodynamics (MHD) origin of viscosity and its decay after the superoutburst (Osaki et al. 2001).

Kato et al. (1998) noted that systems with rebrightening(s) and systems with positive P_{dot} are well correlated. As introduced in subsection 6.2, Kato et al. (1998) considered that positive P_{dot} is a result of expansion of the disk beyond the 3:1 resonance, and suggested an interpretation that the matter beyond the 3:1 resonance produces rebrightenings. Although such an expansion of the disk has not been confirmed by numerical simulations (it is essentially difficult to introduce disk instability-type condition in 2-dimensional or 3-dimensional hydrodynamic simulation), this “reservoir of cool matter” beyond the 3:1 resonance has been favored by infrared excess (subsection 4.6) and the presence of Na D absorption during the rebrightening phase (subsection 4.7). We consider the matter beyond the 3:1 resonance is still a viable hypothesis. Osaki et al. (2001) also followed this interpretation as the source of mass supply to the disk.

In the original TTI model (Osaki 1989), the termination of the superoutburst could not be physically derived but was treated as a free parameter to best reproduce the observation. It was naively (with some observational support) considered that when the disk no longer becomes sufficiently eccentric, the tidal instability became insufficient to sustain the superoutburst and the rapid decline starts.

Hellier (2001b) reconsidered this issue and proposed that in objects with small tidal torques (i.e. small- q systems with $q \lesssim 0.07$) the state of the superoutburst can be quenched even in the presence of an eccentric disk. Hellier (2001b) showed this decoupling of thermal and tidal instabilities can explain both rebrightenings in WZ Sge-type objects and short recurrence times of ER UMa-type objects (Kato, Kunjaya 1995; Robertson et al. 1995). According to Hellier (2001b), the low- q condition is common between these objects and the only difference is that WZ Sge-type objects have very low mass-transfer rates while ER UMa-type objects show high ones. This idea of decoupling was an extension of the working model by Osaki (1995b) to reproduce the extremely short (19 d) supercycle of RZ LMi, and the smaller tidal effect for the matter beyond the 3:1 resonance followed Kato et al. (1998).

The recently published paper by Meyer, Meyer-Hofmeister (2015) followed the same line as in Osaki et al. (2001) and indicated that rebrightenings can be understood as “repeated reflections of transition waves which mediate changes between the hot and the cool state of the accretion disk and travel back and forth in the outer disk region”. In this interpretation, the inner part of the disk remains permanently hot during the rebrightening phase. Meyer, Meyer-Hofmeister (2015) also explained “mini-rebrightenings” as a result of the intermediate state in the so-called S-shape of the thermal equilibrium curve. Meyer, Meyer-Hofmeister (2015) also supposed the necessity of continuous mass inflow from the outermost region, and they ascribed this source to the storage of matter beyond the 3:1 resonance, as originally suggested by Kato et al. (1998). Meyer, Meyer-Hofmeister (2015) also noted that observations indicate faster rise than decline in re-

brightenings (cf. subsection 4.4), indicating that the heating wave should move faster than the cooling wave.

As the term “echo outbursts” was originally introduced to describe phenomenon in X-ray binaries (Augusteijn et al. 1993), which is somewhat different (repeated upward deviations from the exponential declines in classical X-ray transients) from the phenomenon in WZ Sge-type dwarf novae and these “echoes” in X-ray binaries were interpreted as the irradiation-induced mass-transfer, it was quite natural that there were MTB-type explanations for WZ Sge-type rebrightening. [We should note that Kuulkers et al. (1996) already indicated that the irradiation effect is expected to be much smaller in dwarf novae than in X-ray transients. See also Kuulkers (1998) and Kuulkers (2000)]. Hameury et al. (2000) tried to reproduce rebrightenings in EG Cnc by an enhanced mass-transfer. Hameury et al. (2000) partly succeeded in reproducing the light curve. Patterson et al. (2002) supported this interpretation by the claimed detection of enhanced orbital humps during the rebrightening phase. As is evident from the result of numerical simulation by Hameury et al. (2000), this type of enhanced mass-transfer results a temporary shrinkage of the disk radius (cf. Ichikawa, Osaki 1992). Recent observations of systems with multiple rebrightenings did not show such a shrinkage of the disk radius as determined from the superhump periods with the modern interpretation of precession rates, since the pressure effect can be neglected in cool, post-superoutburst disks (C. Nakata et al. in preparation). We consider that the enhanced mass-transfer model is not currently a viable one for explaining (at least for) repeated rebrightenings.

There are, however, remaining problems to be solved. The above models have no interpretation why some WZ Sge-type objects show rebrightening(s) while others do not. As we have seen in this paper (subsection 7.6), higher- q objects tend to show single rebrightenings. Intermediately low- q objects (around the period minimum in CV evolution) tends to show no rebrightenings. Even lower- q objects show long rebrightenings or repeated rebrightenings (we should note that some higher- q object also show repeated rebrightenings). This sequence of rebrightening type needs to be reproduced by further theoretical studies. The existence of a precursor outburst in long rebrightening and renewed growth of superhumps (subsection 4.4) is also a problem to be solved.

8.4. Intermediate Polar

Since various outburst models (particularly models with standard α_C) assume the magnetism of the white dwarfs, we place this subsection here. Up to now, the only object with confirmed intermediate polar (IP)-type nature is V455 And (Araujo-Betancor et al. 2005a; Silvestri et al. 2012; Bloemen et al. 2013) which has a spin period of 67 s. Various authors (e.g. Warner, Pretorius 2008) imply that the 27.87s oscillation of WZ Sge (in quiescence, e.g. Patterson et al. 1998) may be the spin period of the white dwarf. If it is the case, this is the fastest rotation white dwarf among CVs. There is, however, no direct

confirmation of the IP nature (such as by UV pulsation). There have been arguments about the nature of short-period oscillations in WZ Sge (see e.g. Skidmore et al. 1999; Knigge et al. 2002), which have made the origin of the 27.87-s oscillation still an open question.

Observational evidence for the IP nature in WZ Sge-type objects in general has been weak. Most of them are not strong X-ray emitters, nor the quiescent spectra show lines usually associated with IPs (such as HeII). We should consider that the behavior of most of WZ Sge-type objects needs to be explained without strong magnetism of the white dwarf. There remains a possibility that the presence of the magnetism in certain systems may strengthen the WZ Sge-type feature, such as the long recurrence time.

8.5. Long-Term Trend in Quiescence

Kuulkers et al. (2011) studied long-term trends in WZ Sge using historical photographic, visual and photoelectric (including CCD) data in quiescence and suggested the gradual fading trend between the superoutbursts. Kuulkers et al. (2011) proposed this fading can be explained by the formation of a hole in the disk and gave the consequences in relation to the outburst models and the appearance of the supposed spin period. We should note, however, such an analysis based on visual observations is very dangerous, particularly for WZ Sge. WZ Sge has a notorious close companion which is very difficult to resolve by visual observations. It is most likely many AAVSO observers in the past reported the combined light, resulting brighter magnitudes. This effect is stronger for smaller telescopes, and the secular decline may simply be a reflection of the increase of the telescope size. There was also a “psychological” effect: in the 1980s, the quiescent magnitude of WZ Sge was believed to be around 15.0 (the source is currently unknown) by amateur observers, and many observers reported similar magnitudes, including the VSOLJ observers. This belief should have strongly biased the mean magnitudes. Such psychological biases are frequently present in visual observations – the observed minimum magnitudes were often close to the cataloged magnitudes at the time. Special attention is necessary to use visual observations (especially faint ones) in statistical study.⁹

9. Other Problems

9.1. Long-Period Superhumps in Quiescence

Abbott et al. (1992) detected periodic modulations (89.6 min = 0.0622 d) in AL Com in quiescence. Although

⁹ A good example is SS Aur, whose minimum magnitude was believed to be (or shown in variable star charts) 14.5–15.0 in the 1980s by amateur observers, but is now recognized to be around $V=16.0$. There was secular fading trend in the AAVSO data corresponding to this change, and it is difficult to see whether it is a real change or a psychological effect (cf. the “SS Aur problem”, vsnet-chat 6148). There was a so-called “telephone effect” that some observers communicated by phone before making reports, and the resultant magnitudes of these observers tended to become similar values. The same thing could easily happen if observers see other observers’ quiescent observations.

it was originally suspected to be the orbital period, the true orbital period was later identified, giving an ϵ of 0.098. Patterson et al. (1996) suggested such a large ϵ could arise from the disk close to the 2:1 resonance. There have been similar, but less confirmed, cases: BW Scl (Uthas et al. 2012), EQ Lyn (Szkody et al. 2010, not included in table 6 since no outburst has been recorded) and V455 And (Araujo-Betancor et al. 2005a). The possibility of the disk precession around the radius of 2:1 resonance was examined in Kato, Osaki (2013b) and indeed the observed ϵ can be reproduced with a large, but still reasonable, disk if the disk sufficiently expands in quiescence. It is not known whether the development of the 2:1 resonance in quiescence produces $m=2$ (spiral) pattern as in early superhumps and suppress the $m=1$ (one-armed) mode required for singly peaked superhumps.

9.2. Double-Wave Modulations in Quiescence

It has been well known that quiescent orbital variations of high-inclination WZ Sge-type objects (and SU UMa-type dwarf novae with low mass-transfer rates) frequently have double-wave modulations. Good examples are seen in WZ Sge and AL Com (Patterson et al. 1996), V455 And (Araujo-Betancor et al. 2005a; Kato et al. 2009a), V386 Ser (Mukadam et al. 2010, not included in table 6 since no outburst has been recorded), EZ Lyn (Kato et al. 2009b; Zharikov et al. 2013), BW Scl (Augusteijn, Wisotzki 1997; Kato et al. 2013a). The classically accepted interpretation of this phenomenon is a result of a semi-transparent accretion disk (due to the low mass-transfer rate) which allows the light from the hot spot to escape in two directions (Skidmore et al. 2000).

Zharikov et al. (2008) suggested an interpretation that the disk can reach the 2:1 resonance to produce these double-wave modulations, although Zharikov et al. (2006) initially considered that in quiescence there is no chance that the disc extends as far as the 2:1 resonance.

Kononov et al. (2015) recently proposed another mechanism based on the results of three-dimensional simulations of the gas dynamics. Kononov et al. (2015) suggested that the interaction between a precessing spiral density wave, which is essentially at rest in the observer’s frame, and shock regions produces enhances the energy release and is observed as humps. This interpretation can explain the occasional presence of four humps in one orbital cycle.

9.3. Mini-brightening in Quiescence

Szkody et al. (2006) reported temporary brightening of EZ Lyn in quiescence by 0.5 mag. The phenomenon lasted at least a few hours. Zharikov et al. (2006) reported similar phenomena with durations of 8–12 hr in V406 Vir (a WZ Sge-type candidate which has not undergone an outburst). Both authors favored variation in the mass-transfer as the cause of such phenomena. Aviles et al. (2010) further reported a period of 9.28 hr for this phenomenon in V406 Vir. Aviles et al. (2010) suggested the 2:1 resonance as the origin of double wave modulations in quiescence and variation in the mass-transfer as the cause of mini-brightening in quiescence. Aviles et al.

(2010) speculated some kind of feedback between the disk and the secondary to explain the recurrent nature of mini-brightening. Currently only two objects are known to show this type of variation.

10. List of WZ Sge-Type Dwarf Novae

10.1. List of Confirmed Objects

We present an updated list of WZ Sge-type dwarf novae, approximately following the style of table 5 in Kato et al. (2001). The list covers the object up to Kato et al. (2015), approximately up to 2015 January. We included objects having properties (almost) unique to WZ Sge-type dwarf novae: existence of early superhumps (or long delay in the appearance of ordinary superhumps) and/or multiple rebrightenings. A few object were selected based on the outburst properties (large outburst amplitude and the lack of previous outbursts) and other circumstantial evidence (written in remarks). Candidate period bouncers which showed superoutbursts were also included. We only included objects in which early superhumps or ordinary superhumps were detected. For the three very well-known objects (WZ Sge, AL Com and GW Lib), we restricted the references to basic or representative ones (we mostly refer to papers on optical observations and did not include those on detailed observations in quiescence or on theoretical considerations). For other objects, we listed more complete references. The table is in the approximate order of recognition as a WZ Sge-type member, following the table format in Kato et al. (2001) (this agrees to the epoch of outburst for recent objects).

The maximum and minimum magnitudes refer to the V -band or SDSS g' -band whenever available. In other cases, we used other bands (such as R , photographic, unfiltered CCD magnitudes). Since most WZ Sge-type dwarf novae have color indices in outburst close to 0 from V to I , these magnitudes are good approximations to the V -magnitudes. When the true outburst maximum was apparently missed for a long time (several days or more), we supplied the “]” mark (lower limit) for the maximum magnitude. Although most of actual data are indeed lower limits (true maxima are difficult to catch), we did not supply the mark of lower limit for the objects whose early part of the outburst was reasonably observed (e.g. sufficient detection of early superhumps). Whenever available, we adopted modern SDSS g' -magnitudes for the minimum (c.f. Kato et al. 2012b). When modern CCD magnitudes are not available, we used magnitudes from plate scans. These magnitudes are often uncertain due to the faintness of the objects in quiescence. We supplied “:” (uncertain) sign for such values.

The values of P_{SH} mostly refer to stage B superhumps. This is not a serious issue since most WZ Sge-type dwarf novae show only stage B superhumps and the duration of stage C is short, if present.

The years of outbursts do not include likely normal outbursts (i.e. faint outbursts, which are usually single plate detections). The information for these outbursts is given as remarks.

QY Per may belong to a class of long-period WZ Sge-type systems (Kato et al. 2009a). No early superhumps have yet been detected in this system.

10.2. Excluded Objects

One object, which has been listed as WZ Sge-type objects in some catalogs (like AAVSO VSX) have been excluded from the table.

V453 Nor: Although Imada, Monard (2006) reported the possible detection of early superhumps during the 2005 outburst, this part of the light curve resemble the fading part of a precursor outburst of an ordinary SU UMa-type dwarf nova. The variation looks rather irregular and stage A superhumps may have mimicked double-wave modulations of early superhumps. Two further normal outbursts were detected and Soejima et al. (2009) suggested that this object may not be a very typical WZ Sge-type dwarf nova. There was another superoutburst in 2014. Although stage A superhumps were observed, early superhumps were not observed (Kato et al. 2015). The range of variability is 12.6–17.1 (V). The outburst amplitude of 4.5 mag is also small for a WZ Sge-type dwarf nova.

10.3. Current Status of Candidates in Kato et al. (2001)

The updated information of the candidates listed in Kato et al. (2001) are as follows (other than already confirmed as WZ Sge-type dwarf novae and listed in table 6).

RY Dor: Although there is no spectroscopic data, the object (LMCN 1926-09a) is classified as a nova with a slow decline of $t_3=200$ d and the maximum absolute magnitude of -6.9 in Shafter (2013). This object is more likely a slow nova in the LMC.

KY Ara: Schaefer, Hoffleit (1994) suggested either a large-amplitude dwarf nova, a fast nova in the SMC or a gravitational microlensing event.

V359 Cen: The object is a well-confirmed ordinary SU UMa-type dwarf nova (Kato et al. 2002a; Woudt, Warner 2001; Kato et al. 2009a; Kato et al. 2014a).

KX Aql: The object is a well-confirmed ordinary SU UMa-type dwarf nova with relatively rare outbursts Kato et al. (2010).

V336 Per: The object is a large-amplitude dwarf nova without superhumps (see a discussion in Kato et al. 2012b).

IO Del: There was one confirmed outburst at an unfiltered CCD magnitude of 16.73 on 2012 December 17. The astrometry of the object is in agreement with the cataloged object (vsnet-alert 15194).

AP Cru: Woudt, Warner (2002b) detected an orbital period of 5.12 hr and a stable modulation at 1837 s. The object is most likely an intermediate polar and the 1936 outburst was most likely a nova eruption.

CI Gem: The object is an ordinary SU UMa-type dwarf nova with relatively rare outbursts Kato et al. (2009a).

NSV 895: The object is most likely a supernova with $M_{\text{pg}} = -19$ in UGC 2172 at a distance of 1 Mpc (Kato et al. 2012b).

AO Oct: The object is a well-confirmed ordinary SU UMa-type dwarf nova (Patterson et al. 2003; Woudt et al. 2004; Kato et al. 2014a; Mason, Howell 2003).

V551 Sgr: The object is a well-confirmed ordinary SU UMa-type dwarf nova (Kato et al. 2009a; Mason, Howell 2003).

GO Com: The object is a well-confirmed ordinary SU UMa-type dwarf nova (Imada et al. 2005; Kato et al. 2009a; Kato et al. 2010; Kato et al. 2013a).

10.4. AM CVn-Type Objects with Multiple Rebrightenings

Up to now, no definite detection of early superhumps in AM CVn-type objects has been reported, although these systems are expected to have low mass-ratios comparable to hydrogen-rich WZ Sge-type objects [for recent reviews of AM CVn-type objects, see e.g. Nelemans (2005); Solheim (2010)].

Quite recently, SDSS J090221.35+381941.9 showed slow growth of superhumps similar to (hydrogen-rich) period bouncers. This superoutburst was preceded by a separate precursor and followed by a rebrightening. This object may be analogous to WZ Sge-type objects except that it did not show early superhumps (Kato et al. 2014c).

Two recent AM CVn-type objects displayed multiple rebrightenings: ASASSN-14ei (Prieto et al. 2014b; Prieto et al. 2014a; at least 12 times) and ASASSN-14mv (cf. Denisenko et al. 2014; at least 10 times). According to Denisenko et al. (2014), ASASSN-14mv underwent outbursts in 1938 and 2011. The 2011 outburst was also a superoutburst, and the interval between the recent two superoutbursts ~ 1430 d. Although these objects may be helium counterpart of WZ Sge-type objects, we keep it an open question since it is not yet known whether the physics of the rebrightening is the same as hydrogen-rich systems. AM CVn-type objects tend to more complex light curves of superoutbursts (Kato et al. 2013a; Kato et al. 2014b; Levitan et al. 2015). Oscillations similar to WZ Sge-type objects were observed in objects with frequent outbursts (V803 Cen: Kato et al. 2004b; V406 Hya: Nogami et al. 2004), and this behavior may not be unique to helium dwarf novae with very infrequent outbursts. Kato et al. (2004b) suggested a possibility that the helium disk is more difficult to maintain the hot state and transitions to the cool state may occur more frequently than in hydrogen disks. This possibility needs to be explored by theoretical modeling, which may provide a clue to understanding the observed phenomena in hydrogen disks, too.

11. Summary

We have summarized the current understanding and recently obtained findings about WZ Sge-type dwarf novae. We also reviewed the historical development of the understanding of these objects, provided the modern criteria, and reviewed the past research in relation to superhumps, early superhumps and the outburst mechanism.

We provided the updated list of nearly 100 WZ Sge-

Table 6. Confirmed WZ Sge-type dwarf novae.

Object	Max	Min	Amp*	P_{orb}^{\dagger}	P_{SH}^{\ddagger}	Outbursts §	Remarks $^{\parallel}$	References
WZ Sge	7.0	15.3	8.3	0.05669	0.05722	1913, 1946, 1978, 2011	A/B, 1, 2	1, 2, 3, 4, 5, 6, 7, 8
AL Com	12.8	19.8	7.0	0.05667	0.05722	1961, 1965, 1975, 1995, 2001, 2007, 2013	A/B, 3	2, 8, 9, 10, 11, 12, 13, 14, 15
EG Cnc	11.9	18.0	6.1	0.05997	0.06034	1977, 1996	B, 4, 5	16, 17, 18, 19
V2176 Cyg	13.3	19.9	6.6	–	0.0561	1997	A	20, 21, 22, 23
HV Vir	11.2	19.2	8.0	0.05707	0.05820	1929, 1970, 1992, 2002, 2008	D/C?, 6	8, 24, 25, 26, 27, 28, 29, 30, 31
RZ Leo	12.1	18.7	6.6	0.07603	0.07853	1918, 1984, 2000, 2006	C, 4, 7	8, 31, 32, 33, 34, 35, 36
QZ Lib	11.2	18.8	7.6	–	0.06460	2004	B, 4	8, 37
UZ Boo	11.5	19.7	8.2	–	0.0620	1929, 1937, 1938, 1978, 1994, 2003, 2013	B, 8	8, 15, 38, 39, 40
V592 Her	12.3	21.4	9.1	0.0561	0.05661	1968, 1998, 2010	D, 6, 9	8, 39, 41, 42, 43, 44, 45, 46, 47, 48
UW Tri	14.6	22.4	7.8	0.05334	0.05419	1983, 1995, 2008	D, 9	8, 49, 50, 51
CG CMa	13.7	[20]6.3	–	0.0636	1934, 1999	A	8, 52, 53, 54
LL And	12.6	20.1	7.5	0.05506	0.05658	1979, 1993, 2004	–	8, 31, 55, 56
V1108 Her	12.0	17.1:	5.1:	0.05672	0.05748	1932, 1934, 2004	D, 10	8, 57, 58, 59
FL Psc	10.5	17.5	7.0	0.05610	0.05709	1938, 2004	C, 6, 11	8, 60, 61, 62
V498 Vul	15.6	22.5	6.9	–	0.05990	2005, 2008	–, 12	8
DV Dra	15.0	22.2:	7.2:	0.05883	–	1984, 2005	–, 9	8, 63, 64, 65
V572 And]16.4	[22.0]5.6	0.05487	0.05554	2005	A, 9, 13	8, 66, 67
HO Cet	12.0	19.0:	7.0:	0.05490	0.05599	2006	–, 9,	8
ASAS J102522	12.2	19.3	7.1	0.06136	0.06337	2006, 2011	C, 4, 6, 9	8, 62, 68

*Amplitude of outburst (mag).

 † Orbital period (d). ‡ Superhump period (d). § (Likely) normal outbursts are not listed. $^{\parallel}$ Rebrightening type and remarks.

type dwarf novae mainly based on the data obtained by the VSNET Collaboration up to Kato et al. (2015) and discussed the statistics.

The major findings we obtained can be summarized as follows.

- WZ Sge-type dwarf novae are best defined as objects showing early superhumps. These variations are considered to be manifestation of the 2:1 resonance, and the resonance condition makes a distinction between WZ Sge-type dwarf novae and ordinary SU UMA-type dwarf novae. In addition to this, the presence of long or multiple rebrightenings is also (almost) unique to this class of objects.
- The median outburst amplitude is 7.7 mag and the majority of objects have amplitudes larger than 7.0 mag.
- The orbital periods are mostly below 0.06 d, but there are objects with longer periods, some of which are considered to be period bouncers.
- The median interval of observed superoutbursts is 11.5 yr, although this statistics is affected by various biases. The shortest known interval of superoutbursts is 450 d.

- We reviewed the outburst type and showed representative examples of the light curves: type-A outbursts (long-duration rebrightening), type-B outbursts (multiple rebrightenings), type-C outbursts (single rebrightening), type-D outbursts (no rebrightening), and type-E outbursts (double superoutbursts).
- The outburst type is generally reproduced in each object on different superoutburst occasions, but there was an apparent exception in WZ Sge in 1946.
- We presented the updated atlas of early superhumps.
- Early superhumps are shown to grow during the rising phase of the superoutburst.
- The amplitudes of early superhumps systematically decay as the outburst proceeds. The decay is slower in magnitude scale than flux scale.
- The mean amplitudes of early superhumps are mostly less than 0.1 mag, and only a few systems show amplitudes larger than 0.2 mag. We can, however, detect early superhumps with amplitude larger than 0.02 mag in 63% of the studied WZ Sge-type dwarf novae, making early superhumps a useful distinguishing feature for WZ Sge-type dwarf novae.

Table 6. Confirmed WZ Sge-type dwarf novae (continued).

Object	Max	Min	Amp*	P_{orb}^{\dagger}	P_{SH}^{\ddagger}	Outbursts §	Remarks $^{\parallel}$	References
EZ Lyn	12.0	17.8	5.8	0.05901	0.05954	2006, 2010	B, 2	8, 59, 62, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81
IK Leo	13.9	20.7	6.8	–	0.05631	2006	A, 6	8, 82, 83, 84, 85
SS LMi	15.7	21.9	6.2	0.05664	–	1980, 2006	–, 9, 14	8, 86, 87
GW Lib	8.3	17.2	8.9	0.05332	0.05473	1983, 2007	D, 15	8, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99
V455 And	8.6	16.1	7.5	0.05631	0.05713	2007	D, 2	8, 100, 101, 102, 103, 104, 105, 106
1RXS J023238	10.9	18.2	7.3	–	0.06617	2007	B	8
KK Cnc	12.3	20.7	8.4	–	0.06105	2007	D?, 6, 16	8
OT J111217	11.5	20.9	9.4	0.05847	0.05897	2007	D?	8
DY CMi	11.4	19.4	8.0	–	0.06074	2008	B	8, 107, 108, 109
CRTS J090239	16.0	23.2	7.2	0.05652	–	2008	–, 6, 9	8, 110
V466 And	12.8	21.2:	8.4:	0.05637	0.05720	2008	D, 9	8, 111, 112, 113
CT Tri	14.3	21.7:	7.4:	0.05281	0.05366	2008	–, 6, 9	8, 114, 115
V358 Lyr	15.9	[23.2]7.3	–	0.05563	1965, 2008	A, 17	8, 116, 117, 118, 119
CRTS J223003	14.4	21.0:	6.6:	0.05841	–	2009	–, 9	48
SDSS J161027	13.9	20.1	6.2	0.05687	0.05782	1998, 2009	–, 18	48, 120, 121
VX For	12.2	20.6:	8.4:	–	0.06133	1990, 2009	B, 6	48, 122, 123
OT J213806	8.4	16.3	7.9	0.05452	0.05502	1942, 2010, 2014	D, 6, 19	48, 62, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133
EL UMa	13.7	20.3	6.6	–	0.06045:	1981, 2003, 2009	B, 20	48, 134
CRTS J104411]12.6	19.3]6.7	0.05909	0.06024	2010	C, 6, 9	48

*Amplitude of outburst (mag).

 \dagger Orbital period (d). \ddagger Superhump period (d). \S (Likely) normal outbursts are not listed. \parallel Rebrightening type and remarks.

- We computed theoretical light curves assuming the model by Uemura et al. (2012). The result supports that the existence of many objects with low-amplitude early superhumps is consistent with low inclinations of these systems. The model predicts a larger number of high-amplitude systems than observation, which needs to be resolved.
- As reported in earlier works, the outburst type has a strong correlation with period variation of superhumps and the orbital period.
- The delay of appearance of ordinary superhumps is strongly correlated with the orbital period. We consider this is a result of the stronger 2:1 resonance in short-period systems.
- Using the recently developed method of measuring mass ratios using developing phase of superhumps (stage A superhumps), we have been able to measure mass ratios of many WZ Sge-type objects. Mass ratios of WZ Sge-type object have a peak around 0.07–0.08.
- We have identified a limit of mass ratio enabling the 2:1 resonance to be 0.09.
- By using these mass ratios, we showed that there is a linear relation between the period variation of superhumps and the mass ratio in WZ Sge-type objects. By using this relation, we were able to draw an evolutionary picture of a large number of WZ Sge-type.
- We have identified the type of outburst to be an evolutionary sequence: type C \rightarrow D \rightarrow A \rightarrow B \rightarrow E, with some outliers for type-B objects.
- The duration of stage A (evolutionary phase) of superhumps is also well correlated with the estimated mass ratios, supporting that this duration reflects the growth time of the 3:1 resonance.
- By using these new tools (mass ratios from stage A superhumps) and duration of stage A, we have been able to identify candidates for period bouncers better than any existing criteria. Combined with mass ratios from period variation of superhumps, some of the objects with multiple rebrightenings are considered to be period bouncers.

Table 6. Confirmed WZ Sge-type dwarf novae (continued).

Object	Max	Min	Amp*	P_{orb}^{\dagger}	P_{SH}^{\ddagger}	Outbursts §	Remarks $^{\parallel}$	References
SDSS J160501	12.0	19.8	7.8	0.05666:	–	2010	–, 9	62
OT J012059	12.3	20.1	7.8	0.05716	0.05783	2010	A/B, 9	62
PT And	15.8	22	6.2:	–	0.056	1957, 1983, 1986, 1988, 1998, 2010	–, 21	136, 137, 138, 139
OT J230425]13.7	21.1]7.4	–	0.06628	2010	–, 22	62, 140, 141
MisV1443	12.8	20.5:	7.7:	–	0.05673	2011	C	62
V355 UMa	9.9	17.7	7.8	0.05729	0.05809	2011	D	62, 142, 143
OT J210950]11.5	18.7]7.2	0.05865	0.06005	2011	D, 6, 23	144, 145
SDSS J220553]14.4	20.1]5.7	0.05752	0.05815	2011	–, 24	144, 146, 147, 148, 149
SV Ari	14.0:	22.1	8.1:	–	0.05552	1905, 2011	D, 6, 25	144, 150, 151, 152
OT J184228	11.8	20.6:	8.8:	0.07168	0.07234	2011	E+C, 26	144, 153, 154, 155
BW Scl	9.6	16.5	6.9	0.05432	0.05500	2011	D	144, 156, 157, 158, 159
PR Her	12.9	21.0	8.1	0.05422	0.05502	1949, 2011	–,	144, 160
TCP J061128]15.8	[21.0]6.2	0.056:	–	2011	–, 27	–
CRTS J055721]14.7	21.0:]6.3	–	0.05976	2011	–	144
CRTS J001952]15.6	21.5:]5.9	–	0.05677	2012	–	144
MASTER J211258	14.1	21.3	7.2	0.05973	0.06023	2012	B, 9	161
SSS J224739	11.0	20.5:	9.5:	–	0.05667	2006, 2012	B:, 28	162
OT J232727	13.9	21.8	7.9	0.05277	0.05344	2012	–, 9	162, 163, 164
MASTER J203749	14.1	21.3:	7.2:	0.06062	0.06131	2012	B, 9	161
MASTER J081110]14.1	22.1]8.0	–	0.05814	2012	–, 29	162, 165
SSS J122221]11.8	18.8]7.0	–	0.07649	2013	E?, 30	166, 167, 168, 169, 170, 171
OT J112619]14.8	21.8]7.0	0.05423	0.05489	2013	–, 9	162

*Amplitude of outburst (mag).

 † Orbital period (d). ‡ Superhump period (d). § (Likely) normal outbursts are not listed. $^{\parallel}$ Rebrightening type and remarks.

- We have shown that the magnitude when ordinary superhumps appear can be used as a standard candle and have shown that many WZ Sge-type objects have relatively homogeneous quiescent absolute magnitudes. Candidate period bouncers have slightly fainter (by 0.8 mag) quiescent absolute magnitudes.

Acknowledgements

This work was supported by the Grant-in-Aid “Initiative for High-Dimensional Data-Driven Science through Deepening of Sparse Modeling” (25120007) from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan. We are grateful to many observers who have contributed vital observations to the VSNET Collaboration and the VSOLJ database. We acknowledge with thanks the variable star observations from the AAVSO International Database contributed by observers worldwide and used in this research. The survey papers leading to this work were particularly indebted to the Catalina Real-time Transient Survey, MASTER net-

work and ASAS-SN team for making their real-time detection of transient objects available to the public. This work is inspired by many collaborators who have contributed to a series of “Pdot” papers, and the we are grateful for discussions with them. We are grateful to M. Uemura for providing his code to model the light curve of early superhumps.

References

- Abbott, T. M. C., Fleming, T. A., & Pasquini, L. 1997, *A&A*, 318, 134
- Abbott, T. M. C., Robinson, E. L., Hill, G. J., & Haswell, C. A. 1992, *ApJ*, 399, 680
- Alksnis, A., & Zacs, L. 1981, *IBVS*, 1972
- Alksnis, A., & Zharova, A. V. 2000, *IBVS*, 4909
- Antipin, S. V., Samus, N. N., & Kroll, P. 2004, *IBVS*, 5544
- Arai, A. 2010, *Cent. Bur. Electron. Telegrams*, 2275, 7
- Arai, A., & Nogami, D. 2013, *Astron. Telegram*, 4811
- Araujo-Betancor, S., et al. 2005a, *A&A*, 430, 629
- Araujo-Betancor, S., Gänsicke, B. T., Long, K. S., Beuermann, K., de Martino, D., Sion, E. M., & Szkody, P. 2005b, *ApJ*, 622, 589

Table 6. Confirmed WZ Sge-type dwarf novae (continued).

Object	Max	Min	Amp*	P_{orb}^{\dagger}	P_{SH}^{\ddagger}	Outbursts §	Remarks $^{\parallel}$	References
OT J075418	14.7	22.8	8.1	–	0.07076	2013	–, 31	141
TCP J153756	13.6	21.7:	8.1:	0.06101	–	2013	–, 9	162
GR Ori	13.0	22.4	9.4	–	0.05833	1916, 2013	D	162, 172, 173, 174, 175
MASTER J165236	14.8	22.1	7.3	–	0.08473	2013	–, 32	162, 176
OT J062703	12.2	20.3:	8.1:	0.05787	0.05903	2013	–, 9, 33	162
MASTER J181953]13.9	21.6]7.7	0.05684	0.05752	2013	A/B, 9	162, 177
PNV J191501	9.8	18.5	8.7	0.05706	0.05838	2013	D, 6, 9, 34	178, 179, 180
ASASSN-13ax]13.5	21.2]7.7	–	0.05616	2013	A	162, 181
ASASSN-13ck]12.9	20.8]7.9	0.05535	0.05619	2013	A	15
OT J210016]14.2	22.0:]7.8	0.05787	0.05850	2013	–, 9, 35	15
TCP J233822	13.6	21.5	7.9	0.05726	0.05787	2013	B	15
MASTER J061335]14.2	[22.0]7.8	–	0.05609	2013	–, 6, 36	15, 182
MASTER J005740]15.4	20.9]5.5	0.05619	0.05707	2013	–, 2, 6	15, 183
OT J013741]14.5	[20.0]5.5	0.05854	–	2014	–	15
ASASSN-14ac]14.5	21.6]7.1	–	0.05855	2014	–	15
OT J060009]12.6	20.2]6.7	–	0.06331	2014	B	141
MASTER J175924	12.9	21.5:	8.6:	0.05753	0.05810	2014	–, 9 38	184, 185, 186
PNV J171442	10.0	16.8:	6.8:	0.05956	0.06009	2014	B, 9	141
PNV J172929	12.1	21.5	9.4	0.05973	0.06028	2014	D, 40	124
ASASSN-14cl	10.7	18.8	8.1	0.05838	0.06001	2014	D, 6, 9	124, 187, 188
ASASSN-14cq	13.7	21.3:	7.6:	0.05660	0.05735	2014	–, 9	124
ASASSN-14cv	11.2	19.2	8.0	0.05992	0.06041	2014	B, 9	141
FI Cet	14.4	21.6	7.2	0.05594	0.05691	2001, 2014	–,	124, 189
OT J230523	12.3	19.8	7.5	0.05456	0.05560	2014	–, 6, 9	124

*Amplitude of outburst (mag).

 † Orbital period (d). ‡ Superhump period (d). § (Likely) normal outbursts are not listed. $^{\parallel}$ Rebrightening type and remarks.

- Augusteijn, T. 1994, *A&A*, 292, 481
- Augusteijn, T., Kuulkers, E., & Shaham, J. 1993, *A&A*, 279, L13
- Augusteijn, T., & Wisotzki, L. 1997, *A&A*, 324, L57
- Aviles, A., et al. 2010, *ApJ*, 711, 389
- Ayani, K., & Kato, T. 2006, *Cent. Bur. Electron. Telegrams*, 753, 1
- Baba, H., Kato, T., Nogami, D., Hirata, R., Matsumoto, K., & Sadakane, K. 2000, *PASJ*, 52, 429
- Baba, H., et al. 2002, *PASJ*, 54, L7
- Bailey, J. 1979, *MNRAS*, 189, 41P
- Balanutsa, P., et al. 2013, *Astron. Telegram*, 5555
- Balanutsa, P., et al. 2015, *Astron. Telegram*, 6946
- Barwig, H., Mantel, K. H., & Ritter, H. 1992, *A&A*, 266, L5
- Bath, G. T. 1973, *Nature Physical Science*, 246, 84
- Bertola, F. 1964, *Annales d'Astrophysique*, 27, 298
- Bloemen, S., Steeghs, D., De Smedt, K., Vos, J., Gänsicke, B. T., Marsh, T. R., & Rodriguez-Gil, P. 2013, *MNRAS*, 429, 3433
- Boyd, D. 2003, *J. Br. Astron. Assoc.*, 113, 208
- Bullock, E., et al. 2011, *AJ*, 141, 84
- Cannizzo, J. K. 1996, in *IAU Colloq. 158, Cataclysmic Variables and Related Objects*, ed. A. Evans, & J. H. Wood (Dordrecht: Kluwer Academic Publishers), p. 135
- Cannizzo, J. K. 1998, *ApJ*, 493, 426
- Cannizzo, J. K., Lee, H. M., & Goodman, J. 1990, *ApJ*, 351, 38
- Cannizzo, J. K., Still, M. D., Howell, S. B., Wood, M. A., & Smale, A. P. 2010, *ApJ*, 725, 1393
- Chochol, D., Katysheva, N. A., Shugarov, S. Y., & Volkov, I. M. 2009, *Contr. of the Astron. Obs. Skalnaté Pleso*, 39, 43
- Chochol, D., Katysheva, N. A., Shugarov, S. Y., Volkov, I. M., & Andreev, M. V. 2010, *Contr. of the Astron. Obs. Skalnaté Pleso*, 40, 19
- Chochol, D., Katysheva, N. A., Shugarov, S. Y., Zemko, P. O., & Andreev, M. V. 2012, *Contr. of the Astron. Obs. Skalnaté Pleso*, 42, 39
- Christensen, E. J. 2006, *Cent. Bur. Electron. Telegrams*, 746
- Copperwheat, C. M., et al. 2013, *Astron. Telegram*, 5195
- Davis, A. B., Shappee, B. J., Archer Shappee, B., & ASAS-SN 2015, *American Astron. Soc. Meeting Abstracts*, 225, #344.02
- della Valle, M., & Augusteijn, T. 1990, *Messenger*, 61, 41
- Denisenko, D., et al. 2012, *Astron. Telegram*, 4506
- Denisenko, D., et al. 2013a, *Astron. Telegram*, 4881
- Denisenko, D., et al. 2014, *Astron. Telegram*, 6857
- Denisenko, D., et al. 2013b, *Astron. Telegram*, 5399
- Djorgovski, S. G., et al. 2008, *Astron. Telegram*, 1411
- Downes, R. A. 1990, *AJ*, 99, 339
- Downes, R. A., & Margon, B. 1981, *MNRAS*, 197, 35P

Table 6. Confirmed WZ Sge-type dwarf novae (continued).

Object	Max	Min	Amp*	P_{orb}^{\dagger}	P_{SH}^{\ddagger}	Outbursts §	Remarks $^{\parallel}$	References
ASASSN-14gx	14.9	21.7:	6.8:	0.05488	0.05609	2014	-, 9, 41	124
ASASSN-14jf	13.3	21.0:	7.7:	0.05539	0.05595	2014	-, 9	124
OT J030929	11.0	18.9	7.9	0.05615	0.05744	2014	D, 6, 9	124
ASASSN-14jq]13.7	20.5]6.8	-	0.05518	2014	A, 42	124
ASASSN-14jv	11.3	19.3	8.0	0.05442	0.05510	2014	D, 9	124
ASASSN-14mc	14.3	21.0:	6.7:	-	0.05546	2014	-, 43	124
ASASSN-15ah	13.6	21.8:	8.2:	-	0.05547	2015	-, 44	124
ASASSN-15bp	11.9	20.5	8.6	0.05563	0.05670	2014	B?	124, 190
MASTER J085854]13.7	18.6:]4.9	-	0.05556	2009?, 2014	B, 45	124, 191
Objects in Mroz et al. (2013):								
OGLE-GD-DN-001	17.8	[21.6]3.8	-	0.06072	2007	B	192
OGLE-GD-DN-014]18.4	22.6]4.2	-	0.08931	2006	B, 46	192
Borderline or long-period systems:								
V1251 Cyg	12.5	20.5	8.0	0.07433	0.07597	1963, 1991, 1994, 1997, 2008	C, 9, 47	8, 193, 194, 195
BC UMa	10.9	18.5	7.6	0.06261	0.06457	1960, 1962, 1982, 1990, 1992, 1994, 1995, 2000, 2003, 2009	C, 6, 48	8, 196, 197, 198, 199, 200
MASTER J004527]12.5	19.3:]6.8	-	0.08037	2013	C, 49	15, 201
EI Psc-type objects:								
CSS J174033	12.8	19.5:	6.7:	0.04505	0.04559	2007, 2013, 2014	A, 6, 9, 50	162, 124, 202, 203, 204

*Amplitude of outburst (mag).

 † Orbital period (d). ‡ Superhump period (d). § (Likely) normal outbursts are not listed. $^{\parallel}$ Rebrightening type and remarks.

- Drake, A. J., et al. 2009, ApJ, 696, 870
Drake, A. J., et al. 2013, Astron. Telegram, 4699
Duerbeck, H. W. 1984, IBVS, 2502
Duerbeck, H. W. 1987, Space Sci. Rev., 45, 1
Duerbeck, H. W., & Mennickent, R. E. 1998, IBVS, 4637
Duerbeck, H. W., Schmeer, P., Knapen, J. H., & Pollacco, D. 1999, IBVS, 4759
Duerbeck, H. W., & Seitter, W. C. 1987, Ap&SS, 131, 467
Gammie, C. F., & Menou, K. 1998, ApJ, 492, L75
Gänsicke, B. T., et al. 2009, MNRAS, 397, 2170
Gänsicke, B. T., et al. 2006, MNRAS, 365, 969
Gänsicke, B. T., Szkody, P., Howell, S. B., & Sion, E. M. 2005, ApJ, 629, 451
Golovin, A., et al. 2007, IBVS, 5763
Golovin, A., et al. 2005, IBVS, 5611
Golysheva, P., & Shugarov, S. 2014, Contr. of the Astron. Obs. Skalnaté Pleso, 43, 312
Gorbovskey, E. S., et al. 2013, Astron. Rep., 57, 233
Graham, M. L., Broekhoven-Fiene, H., Parker, A. H., Sadavoy, S., Maxwell, A. J., Hsiao, E. Y., & Balam 2010, Cent. Bur. Electron. Telegrams, 2275, 6
Grubissich, C., & Rosino, L. 1958, Asiago Contr., 93, 1
Hōshi, R. 1979, Progress of Theoret. Phys., 61, 1307
Hameury, J.-M., Lasota, J.-P., & Warner, B. 2000, A&A, 353, 244
Harrison, T. E. 1991, IAU Circ., 5233
Harrison, T. E., Hamilton, R. T., Tappert, C., Hoffman, D. I., & Campbell, R. K. 2013, AJ, 145, 19
Harrison, T. E., Johnson, J. J., McArthur, B. E., Benedict, G. F., Szkody, P., Howell, S. B., & Gelino, D. M. 2004, AJ, 127, 460
Hassall, B. J. M. 1985, MNRAS, 216, 335
Hellier, C. 2001a, Cataclysmic Variable Stars: How and why they vary (Berlin: Springer)
Hellier, C. 2001b, PASP, 113, 469
Henden, A., Shears, J., Poyner, G., Dvorak, S., & Krajci, T. 2008, Cent. Bur. Electron. Telegrams, 1582, 1
Hessman, F. V., Robinson, E. L., Nather, R. E., & Zhang, E. 1984, ApJ, 286, 747
Hiroi, K., et al. 2009, PASJ, 61, 697
Hirose, M., & Osaki, Y. 1990, PASJ, 42, 135
Hoffmeister, C. 1951, Erg. Astron. Nachr., 12, 14
Howell, S., & Szkody, P. 1988, PASP, 100, 224
Howell, S. B., De Young, J., Mattei, J. A., Foster, G., Szkody, P., Cannizzo, J. K., Walker, G., & Fierce, E. 1996, AJ, 111, 2367
Howell, S. B., Henden, A. A., Landolt, A. U., & Dain, C. 2004, PASP, 116, 527
Howell, S. B., & Hurst, G. M. 1996, J. Br. Astron. Assoc., 106, 29
Howell, S. B., & Jacoby, G. H. 1986, PASP, 98, 802
Howell, S. B., Mitchell, K. J., & Warnock, A. I. 1987, PASP, 99, 126
Howell, S. B., Nelson, L. A., & Rappaport, S. 2001, ApJ, 550, 897
Howell, S. B., & Szkody, P. 1990, ApJ, 356, 623

Table 6. Confirmed WZ Sge-type dwarf novae (continued).

Remarks: 1. The 1946 superoutburst did not show strong rebrightening comparable to A-type (see text in this paper). The quiescent magnitude is taken from Kuulkers et al. (2011), who reported significant secular variation. 2. Eclipsing system. 3. P_{orb} is taken from the best estimate of the early superhumps in three superoutbursts (Kato et al. 2014a). The 2007 superoutburst showed rebrightenings intermediate between type A and B (Uemura et al. 2008b). The 2015 superoutburst lacked early superhumps. Several normal outbursts have been recorded. For a complete list of outbursts, see Kato et al. (2014a). 4. Early superhumps existed only for a short interval. 5. P_{orb} is taken from Patterson et al. (1998). This period is not in agreement with the period of early superhumps (Matsumoto et al. 1998; Kato et al. 2004a). More observations are desired to resolve this discrepancy [see also discussion in Nakata et al. (2013b)]. There was a well-confirmed normal outburst in 2009 October (Templeton 2009; Lange 2010). 6. Transition to stage C during the superoutburst plateau was recorded. 7. A well-documented normal outburst was recorded in 1989 March (see O’Donoghue et al. (1991)). Richter (1985) recorded several outbursts in archival plates, whose nature has not been determined. P_{orb} has been updated to be 0.0760301(1) d using the CRTS data. The object may better be categorized in long-period systems, but is included in this position since it is one of the most classically identified WZ Sge-type objects. 8. The outbursts before 1978 were taken from Richter (1986a). It is not clear whether the 1937 and 1938 ones were superoutbursts. 9. P_{orb} is taken from the best estimate of the period of early superhumps. 10. The orbital period is from Pavlenko and Antonyuk (in prep.). The quiescent magnitude is uncertain due to the close visual companion. The value is taken from Price et al. (2004). Only two bright historical outbursts in Price et al. (2004) are listed in the table. There were faint (possible) outbursts in 1939 and 1940 (Price et al. 2004). 11. The candidate orbital period is from Kato et al. (2012a). Templeton et al. (2006) reported a spectroscopic orbital period of 0.0569(5) d. 12. May be a borderline object with the minimum recurrence time of ~ 1000 d. Early superhumps have not yet convincingly recorded. During the 2005 outburst, ordinary superhumps were detected 7 d after the outburst maximum, suggesting that the phase of early superhumps was not long. The object is selected as a WZ Sge-type object based on the long duration (more than 20 d) of the superoutburst and nearly zero period derivative (Kato et al. 2009a). 13. The minimum magnitude is from CCD images in our post-outburst images (not plotted in Imada et al. 2006). 14. Superhump variations reported in Shears et al. (2008) were identified as early superhumps in Kato et al. (2009a). There may have been an outburst in 1991 (Harrison 1991), which may be a confusion with a nearby field star. 15. The maximum V -magnitude is taken from Vican et al. (2011). The minimum V -magnitude is from preoutburst AAVSO observations. The object stayed by ~ 1 mag brighter after the 2007 outburst. The historically used value (18.5p) in quiescence appears to be too faint. 16. One-day dip was recorded before the termination of the superoutburst (Kato et al. 2009a). The object was not well observed after the termination of the superoutburst to detect possible rebrightenings. The maximum V -magnitude is taken from ASAS-3 prediscovery observation, which is not reflected on GCVS (Samus et al. 2011) and other catalogs. 17. The upper limit for the quiescent magnitude is from Shears et al. (2010). 18. The 1998 outburst was recorded on one image (15.0 mag, Wils et al. 2010). 19. The object is a fainter companion of a close visual binary. The minimum magnitude is difficult to estimate and is taken from the AAVSO VSX page. Hudec (2010) documented the 1942 outburst. 20. The 1981 and 2003 outbursts refer to single-epoch observations. These outbursts are listed as possible superoutburst based on their brightness. The 2009 outburst was probably detected in the late stage and only rebrightening part was observed [see discussion in Kato et al. (2010)]. The superhump period listed in the table refers to the one recorded during the rebrightenings. 21. Spectroscopically confirmed dwarf nova (A. Arai, vsnet-alert 12528). The most likely superhump period is given (vsnet-alert 12527). Although relatively short recurrence time suggests a system similar to ordinary SU UMa-type dwarf novae, the long duration of outbursts (cf. Alksnis, Zharova 2000) and possible rebrightening(s) support the WZ Sge-type classification. 22. Although no early superhumps were present, C. Nakata et al. in preparation re-classified this object as a likely period bouncer based on the long stage A phase (~ 120 cycles). 23. Suspected orbital period detected from modulations during the outburst. 24. The true maximum was most likely missed for a relatively long period.

- Howell, S. B., Szkody, P., & Cannizzo, J. K. 1995a, *ApJ*, 439, 337
 Howell, S. B., Szkody, P., Kreidl, T. J., Mason, K. O., & Puchnarewicz, E. M. 1990, *PASP*, 102, 758
 Howell, S. B., Szkody, P., Sonneborn, G., Fried, R., Mattei, J., Oliverson, R. J., Ingram, D., & Hurst, G. M. 1995b, *ApJ*, 453, 454
 Howell, S. B., Warnock, A., Mason, K. O., Reichert, G. A., & Kreidl, T. J. 1988, *MNRAS*, 233, 79
 Hu, J.-Y., Qiu, Y.-L., Li, W.-D., Wei, J.-Y., & Esamdin, A. 1997, *IAU Circ.*, 6731
 Hudec, R. 2010, *Astron. Telegram*, 2619
 Huruata, M. 1983, *IBVS*, 2401
 Ichikawa, S., & Osaki, Y. 1992, *PASJ*, 44, 15
 Ichikawa, S., & Osaki, Y. 1994, in *Theory of Accretion Disks-2*, ed. W. J. Duschl, J. Frank, F. Meyer, E. Meyer-Hofmeister, & W. M. Tscharnutter (Dordrecht: Kluwer Academic Publishers), p. 169
 Imada, A., et al. 2005, *PASJ*, 57, 193
 Imada, A., Kubota, K., Kato, T., Nogami, D., Maehara, H., Nakajima, K., Uemura, M., & Ishioka, R. 2006, *PASJ*, 58, L23
 Imada, A., & Monard, L. A. G. B. 2006, *PASJ*, 58, L19
 Ingram, D., & Szkody, P. 1992, *IBVS*, 3810
 Ishioka, R., et al. 2001, *PASJ*, 53, 905
 Ishioka, R., et al. 2003, *PASJ*, 55, 683
 Ishioka, R., et al. 2002, *A&A*, 381, L41
 Isogai, M., Arai, A., Yonehara, A., Kawakita, H., Uemura, M., & Nogami, D. 2015, *PASJ*, 67, 7
 Itagaki, K., et al. 2013, *Cent. Bur. Electron. Telegrams*, 3554, 1
 Kato, T. 1995, *IBVS*, 4152
 Kato, T. 2002, *PASJ*, 54, L11
 Kato, T. 2004, *PASJ*, 56, S135
 Kato, T., et al. 2015, *PASJ*, in press (arXiv/1507.05610)
 Kato, T., et al. 2014a, *PASJ*, 66, 90
 Kato, T., et al. 2013a, *PASJ*, 65, 23
 Kato, T., et al. 2014b, *PASJ*, 66, 30

Table 6. Confirmed WZ Sge-type dwarf novae (continued).

Remarks (continued): 25. The maximum magnitude is from the 1905 observation in Wolf, Wolf (1905) scaled to modern magnitude. The true maximum in 2011 was most likely missed for a relatively long period. There was a possible outburst in 1943 (Duerbeck 1987). 26. One superoutburst with early superhumps and another one with ordinary superhumps. Good candidate for the period bouncer (Kato, Osaki 2013b). One solitary rebrightening was recorded (Katysheva et al. 2013). 27. <<http://www.cbat.eps.harvard.edu/unconf/followups/J06112800+4041087.html>> and vsnet-alert 13871. The period has been tentatively identified as early superhumps based on the description on this page. 28. The 2012 outburst was probably detected in the late stage and only rebrightening part was observed [see discussion in Kato et al. (2014b)]. 29. Likely early superhumps were detected. Most of the phase of early superhumps were likely missed. 30. Double superoutburst resembling OT J184228, although the initial one was not observed in real-time. Good candidate for the period bouncer (Kato et al. 2013b). 31. Although no early superhumps were present, C. Nakata et al. in preparations suggested this object as a period bouncer based on the long stage A phase (~ 190 cycles). 32. Red quiescent counterpart. The long superhump period and the large outburst amplitude suggests a period bouncer. 33. The maximum magnitude is an updated value on the AAVSO VSX page. 34. Although J. Echevarria reported a spectroscopic period of 0.06164 d based on 2.5-hr observation (vsnet-alert 15832), this period is not consistent with the superhump period. The baseline of the observation was probably too short to determine the orbital period. 35. There is no SDSS counterpart. The minimum magnitude is from AASVO VSX page. 36. The evolution of superhumps resembles that of ordinary SU UMA-type dwarf nova. The phase of early superhumps may not have been well recorded (Kato et al. 2014a). 37. The object showed a precursor outburst, rather than early superhumps. The long phase of stage A superhumps, slow decline and multiple rebrightenings suggest that the object is a period bouncer and we classify it as a WZ Sge-type (C. Nakata et al. in preparation). 38. The reported discovery magnitude (12.7 mag) is probably overestimated by 0.5 mag (D. Denisenko, vsnet-alert 17188). The maximum magnitude is taken from Stanek et al. (2014b). The quiescent counterpart is somewhat unclear (vsnet-alert 17188). The superhump period is from an analysis of the VSNET data. 39. Range from AAVSO VSX page. 40. Maximum V -magnitude is from vsnet-alert 17324 (H. Maehara). Candidate period bouncer from the estimated q value (Kato et al. 2015). 41. Possible stage C superhumps. 42. The initial part of the outburst was missed due to the gap in the observation. 43. Included because of delayed appearance of ordinary superhumps and large outburst amplitude. The minimum magnitude is taken from the AAVSO VSX page. 44. Included because of delayed appearance of ordinary superhumps and large outburst amplitude. Likely not an extreme WZ Sge-type object based on the behavior of superhumps (Kato et al. 2015). 45. The possible CRTS detection reported in Balanutsa et al. (2015) was on 2009 October 23 at 14.0 mag. 46. The early part of the outburst was not observed. May be classified a long-period system. 47. During the 2008 superoutburst, a delay in the appearance of ordinary superhumps and likely early superhumps were detected (Kato et al. 2009a). 48. Several possible normal outbursts were recorded, including the well-observed one in 2001. Romano (1964) reported a faint outburst (likely a faint superoutburst) in 1961. The maximum magnitudes of the superoutbursts vary significantly from outburst to outburst. Early superhumps were detected during the 2003 superoutburst (Maehara et al. 2007). 49. Long-period system with a large outburst amplitudes. The phase of early superhumps was probably missed. 50. Object below the period minimum (hydrogen-depleted system). The 2013 and 2014 superoutbursts were both WZ Sge-type ones separated by less than 500 d.

- Kato, T., et al. 2009a, PASJ, 61, S395
 Kato, T., & Kunjaya, C. 1995, PASJ, 47, 163
 Kato, T., et al. 2012a, PASJ, 64, 21
 Kato, T., Maehara, H., & Monard, B. 2008, PASJ, 60, L23
 Kato, T., Maehara, H., & Uemura, M. 2012b, PASJ, 64, 62
 Kato, T., et al. 2010, PASJ, 62, 1525
 Kato, T., Matsumoto, K., Nogami, D., Morikawa, K., & Kiyota, S. 2001a, PASJ, 53, 893
 Kato, T., Matsumoto, K., & Stubbings, R. 1999, IBVS, 4760
 Kato, T., Monard, B., Hamsch, F.-J., Kiyota, S., & Maehara, H. 2013b, PASJ, 65, L11
 Kato, T., Nogami, D., Baba, H., & Matsumoto, K. 1998, in ASP Conf. Ser. 137, Wild Stars in the Old West, ed. S. Howell, E. Kuulkers, & C. Woodward (San Francisco: ASP), p. 9
 Kato, T., Nogami, D., Baba, H., Matsumoto, K., Arimoto, J., Tanabe, K., & Ishikawa, K. 1996, PASJ, 48, L21
 Kato, T., Nogami, D., Lockley, J. J., & Somers, M. 2001b, IBVS, 5116
 Kato, T., Nogami, D., Matsumoto, K., & Baba, H. 2004a, PASJ, 56, S109
 Kato, T., et al. 2014c, PASJ, 66, L7
 Kato, T., & Osaki, Y. 2013a, PASJ, 65, 97
 Kato, T., & Osaki, Y. 2013b, PASJ, 65, 115
 Kato, T., et al. 2009b, PASJ, 61, 601
 Kato, T., Sekine, Y., & Hirata, R. 2001, PASJ, 53, 1191
 Kato, T., Stubbings, R., Monard, B., Butterworth, N. D., Bolt, G., & Richards, T. 2004b, PASJ, 56, S89
 Kato, T., et al. 2002a, A&A, 395, 541
 Kato, T., Uemura, M., Ishioka, R., Nogami, D., Kunjaya, C., Baba, H., & Yamaoka, H. 2004c, PASJ, 56, S1
 Kato, T., Uemura, M., Matsumoto, K., Kinnunen, T., Garrard, G., Masi, G., & Yamaoka, H. 2002b, PASJ, 54, 999
 Katysheva, N., et al. 2013, Central European Astrophys. Bull., 37, 335
 Kimura, M., et al. 2015, PASJ, submitted
 Knigge, C., Baraffe, I., & Patterson, J. 2011, ApJS, 194, 28
 Knigge, C., Hynes, R. I., Steeghs, D., Long, K. S., Araujo-Betancor, S., & Marsh, T. R. 2002, ApJ, 580, L151
 Kolb, U. 1993, A&A, 271, 149
 Kolb, U., & Baraffe, I. 1999, MNRAS, 309, 1034
 Kononov, D. A., Bisikalo, D. V., Puzin, V. B., Zhilkin, A. G., & Sytov, A. Y. 2015, Astron. Rep., 59, 191
 Kunjaya, C., Kato, T., & Hirata, R. 1998, Highlights of Astronomy, 11, 388
 Kurochkin, N. E. 1984, Astron. Tsirk., 1325, 5
 Kuulkers, E. 1998, New Astron. Rev., 42, 1
 Kuulkers, E. 2000, New Astron. Rev., 44, 27
 Kuulkers, E., Henden, A. A., Honeycutt, R. K., Skidmore, W., Waagen, E. O., & Wynn, G. A. 2011, A&A, 528, A152

Table 6. Confirmed WZ Sge-type dwarf novae (continued).

References: 1. Patterson et al. (1981); 2. Ishioka et al. (2002); 3. Kuulkers et al. (2002); 4. Patterson et al. (2002); 5. Baba et al. (2002); 6. Howell et al. (2004); 7. Kato et al. (2008); 8. Kato et al. (2009a); 9. Pych, Olech (1995); 10. Kato et al. (1996); 11. Patterson et al. (1996); 12. Howell et al. (1996); 13. Nogami et al. (1997); 14. Uemura et al. (2008b); 15. Kato et al. (2014a); 16. Huruwata (1983); 17. Matsumoto et al. (1998); 18. Patterson et al. (1998); 19. Kato et al. (2004a); 20. Hu et al. (1997); 21. Vanmunster, Sarneczky (1997); 22. Kwast, Semeniuk (1998); 23. Novák et al. (2001); 24. Duerbeck (1984); 25. Schmeer et al. (1992); 26. Ingram, Szkody (1992); 27. Barwig et al. (1992); 28. Leibowitz et al. (1994); 29. Kato et al. (2001); 30. Ishioka et al. (2003); 31. Patterson et al. (2003); 32. Wolf (1919); 33. Mattei et al. (1985); 34. Richter (1985); 35. Mennickent, Tappert (2001); 36. Ishioka et al. (2001); 37. Schmidtbreick et al. (2004); 38. Bailey (1979); 39. Richter (1992); 40. Kuulkers et al. (1996); 41. Richter (1968); 42. Richter (1991); 43. Duerbeck, Mennickent (1998); 44. van Teeseling et al. (1999); 45. Mennickent et al. (2002); 46. Kato et al. (2002b); 47. Mason, Howell (2003); 48. Kato et al. (2010); 49. Kurochkin (1984); 50. Robertson et al. (2000); 51. Kato et al. (2001b); 52. Duerbeck (1987); 53. Kato et al. (1999); 54. Duerbeck et al. (1999); 55. Wild (1979); 56. Kato (2004); 57. Nakano et al. (2004); 58. Price et al. (2004); 59. Pavlenko et al. (2010); 60. Golovin et al. (2005); 61. Templeton et al. (2006); 62. Kato et al. (2012a); 63. Pavlov, Shugarov (1985); 64. Wenzel (1991a); 65. Richter (1992); 66. Quimby et al. (2005); 67. Imada et al. (2006); 68. Vanmunster et al. (2006); 69. Szkody et al. (2006); 70. Pavlenko et al. (2007); 71. Shears et al. (2007); 72. Zharikov et al. (2008); 73. Pavlenko (2009); 74. Kato et al. (2009b); 75. Pavlenko et al. (2010); 76. Pavlenko et al. (2012); 77. Zharikov et al. (2013); 78. Szkody et al. (2013); 79. Nakata et al. (2013b); 80. Pavlenko et al. (2014); 81. Isogai et al. (2015); 82. Christensen (2006); 83. Ayani, Kato (2006); 84. Golovin et al. (2007); 85. Uemura et al. (2008a); 86. Alksnis, Zacs (1981); 87. Shears et al. (2008); 88. Maza, Gonzalez (1983); 89. Duerbeck, Seitter (1987); 90. Szkody et al. (2000); 91. van Zyl et al. (2000); 92. Thorstensen et al. (2002); 93. Woudt, Warner (2002a); 94. van Zyl et al. (2004); 95. Waagen et al. (2007); 96. Hiroi et al. (2009); 97. Schwieterman et al. (2010); 98. Bullock et al. (2011); 99. Vican et al. (2011); 100. Araujo-Betancor et al. (2005a); 101. Templeton et al. (2007); 102. Kato et al. (2008); 103. Nogami et al. (2009); 104. Maehara et al. (2009); 105. Matsui et al. (2009); 106. Uemura et al. (2012); 107. Yamaoka et al. (2008b); 108. Shears et al. (2009); 109. Woudt, Warner (2011); 110. Djorgovski et al. (2008); 111. Yamaoka et al. (2008a); 112. Shafter et al. (2009); 113. Chochol et al. (2010); 114. Shugarov et al. (2008); 115. Chochol et al. (2009); 116. Richter (1986b); 117. Antipin et al. (2004); 118. Henden et al. (2008); 119. Shears et al. (2010); 120. Wils et al. (2010); 121. Thorstensen, Skinner (2012); 122. Liller et al. (1990); 123. Liller et al. (1999); 124. Kato et al. (2015); 125. Yamaoka (2010); 126. Nakano (2010); 127. Arai (2010); 128. Graham et al. (2010); 129. Tovmassian et al. (2010); 130. Hudec (2010); 131. Chochol et al. (2012); 132. Zemko, Kato (2013); 133. Mitrofanova et al. (2014); 134. Pesch, Sanduleak (1987); 135. Wils et al. (2010); 136. T. Ohshima et al. in preparation; 137. Grubissich, Rosino (1958); 138. Sharov, Alksnis (1989); 139. Zheng et al. (2010); 140. Nakano et al. (2011a); 141. C. Nakata et al. in preparation; 142. Szkody et al. (2006); 143. Gänsicke et al. (2006); 144. Kato et al. (2013a); 145. Yamaoka et al. (2011); 146. Szkody et al. (2003); 147. Warner, Woudt (2004); 148. Szkody et al. (2007); 149. Southworth et al. (2008); 150. Wolf, Wolf (1905); 151. Duerbeck (1987); 152. Robertson et al. (2000); 153. Nakano et al. (2011b); 154. Katysheva et al. (2013); 155. Arai, Osaki (2013b); 156. Augustejn, Wisotzki (1997); 157. Abbott et al. (1997); 158. Gänsicke et al. (2005); 159. Uthas et al. (2012); 160. Hoffmeister (1951); 161. Nakata et al. (2013b); 162. Kato et al. (2014b); 163. Yamaoka et al. (2012); 164. Nakano, Noguchi (2012); 165. Denisenko et al. (2012); 166. Kato et al. (2013b); 167. Drake et al. (2013); 168. Levato et al. (2013); 169. Marsh et al. (2013); 170. Kuulkers et al. (2013); 171. Neustroev, Sjoberg (2013); 172. Thiele (1916); 173. Robertson et al. (2000); 174. Kato et al. (2012b); 175. Arai, Nogami (2013); 176. Denisenko et al. (2013a); 177. Shurpakov et al. (2013); 178. Kato et al. (2014a); 179. Itagaki et al. (2013); 180. Nakata et al. (2013a); 181. Copperwheat et al. (2013); 182. Vladimirov et al. (2013); 183. Balanutsa et al. (2013); 184. Vladimirov et al. (2014); 185. Stanek et al. (2014b); 186. vsnet-alert 17235; 187. Stanek et al. (2014a); 188. Teyssier (2014); 189. Smith et al. (2002); 190. Williams et al. (2015); 191. Balanutsa et al. (2015); 192. Mroz et al. (2013); 193. Weber (1966); 194. Wenzel (1991b); 195. Kato (1995); 196. Romano (1964); 197. Howell et al. (1995b); 198. Kunjaya et al. (1998); 199. Boyd (2003); 200. Maehara et al. (2007); 201. Denisenko et al. (2013b); 202. T. Ohshima et al. in preparation; 203. Prieto et al. (2013); 204. Nesci et al. (2013);

Table 6. Confirmed WZ Sge-type dwarf novae (continued).

Full names of abbreviations: ASAS J102522-1542.4 (ASAS J102522), 1RXS J023238.8-371812 (1RXS J023238), OT J111217.4-353829 (OT J111217), CRTS J090239.7+052501 = CSS080304:090240+052501 (CRTS J090239), CRTS J223003.0-145835 = CSS090727:223003-145835 (CRTS J223003), SDSS J161027.61+090738.4 (SDSS J161027), OT J213806.6+261957 (OT J213806), CRTS J104411.4+211307 = CSS100217:104411+211307 (CRTS J104411), SDSS J160501.35+203056.9 (SDSS J160501), OT J012059.6+325545 (OT J012059), OT J230425.8+062546 (OT J230425), OT J210950.5+134840 (OT J210950), SDSS J220553.98+115553.7 (SDSS J220553), OT J184228.1+483742 (OT J184228), TCP J06112800+4041087 (TCP J061128), CRTS J055721.8-363055 = SSS111229:055722-363055 (CRTS J055721), CRTS J001952.2+433901 = CSS120131:001952+433901 (CRTS J001952), MASTER OT J211258.65+242145.4 (MASTER J211258), SSS J224739.7-362253 = SSS120724:224740-362254 (SSS J224739), OT J232727.2+085539 = PNV J23272715+0855391 (OT J232727), MASTER OT J203749.39+552210.3 (MASTER J203749), MASTER OT J081110.46+660008.5 (MASTER J081110), SSS J122221.7-311523 = SSS130101:122222-311525 (SSS J122221), OT J112619.4+084651 = CSS130106:112619+084651 (OT J112619), OT J075418.7+381225 (OT J075418), TCP J15375685-2440136 (TCP J153756), MASTER OT J165236.22+460513.2 (MASTER J165236), OT J062703.8+395250 = PNV J06270375+3952504 (OT J062703), MASTER OT J181953.76+361356.5 (MASTER J181953), PNV J19150199+0719471 (PNV J191501), OT J210016.0-024258 = CSS130905:210016-024258 (OT J210016), TCP J23382254-2049518 (TCP J233822), MASTER OT J061335.30+395714.7 (MASTER J061335), MASTER OT J005740.99+443101.5 (MASTER J005740), OT J013741.1+220312 = CSS140104:013741+220312 (OT J013741), OT J060009.9+142615 = PNV J06000985+1426152 (OT J060009), MASTER OT J175924.12+252031.7 (MASTER J175924), PNV J17144255-2943481 (PNV J171442), PNV J17292916+0054043 (PNV J172929), OT J230523.1-022546 = PNV J23052314-0225455 (OT J230523), OT J030929.8+263804 = PNV J03093063+2638031 (OT J030929), MASTER OT J085854.16-274030.7 (MASTER J085854), MASTER OT J004527.52+503213.8 (MASTER J004527), CSS J174033.5+414756 = CSS130418:174033+414756 (CSS J174033)

- Kuulkers, E., Howell, S. B., & van Paradijs, J. 1996, *ApJ*, 462, L87
- Kuulkers, E., Knigge, C., Steeghs, D., Wheatley, P. J., & Long, K. S. 2002, in *The Physics of Cataclysmic Variables and Related Objects*, ed. B. T. Gänsicke, K. Beuermann, & K. Reinsch (San Francisco: ASP), p. 443
- Kuulkers, E., Page, K. L., Knigge, C., Marsh, T. R., Osborne, J. P., & Sivakoff, G. R. 2013, *Astron. Telegram*, 4716
- Kwast, T., & Semeniuk, I. 1998, *IBVS*, 4654
- Lange, T. 2010, *BAV Rundbrief*, 59, 65
- Lasota, J. P., Hameury, J. M., & Huré, J. M. 1995, *A&A*, 302, L29
- Leibowitz, E. M., Mendelson, H., Bruch, A., Duerbeck, H. W., Seitter, W. C., & Richter, G. A. 1994, *ApJ*, 421, 771
- Levato, H., et al. 2013, *Astron. Telegram*, 4700
- Levitan, D., Groot, P. J., Prince, T. A., Kulkarni, S. R., Laher, R., Ofek, E. O., Sesar, B., & Surace, J. 2015, *MNRAS*, 446, 391
- Liller, W., Frasca, A., & Messina, S. 1999, *IAU Circ.*, 7327
- Liller, W., Phillips, M., Hamuy, M., Lamontagne, R., Baganoff, F., Maza, J., & Wischnjewsky, M. 1990, *IAU Circ.*, 5127
- Lin, D. N. C., & Papaloizou, J. 1979, *MNRAS*, 186, 799
- Littlefair, S. P., Dhillon, V. S., Marsh, T. R., & Gänsicke, B. T. 2006, *MNRAS*, 371, 1435
- Livio, M., & Pringle, J. E. 1992, *MNRAS*, 259, 23
- Lubow, S. H. 1991a, *ApJ*, 381, 259
- Lubow, S. H. 1991b, *ApJ*, 381, 268
- Maehara, H., Hachisu, I., & Nakajima, K. 2007, *PASJ*, 59, 227
- Maehara, H., et al. 2009, in *ASP Conf. Ser. 404, The Eighth Pacific Rim Conference on Stellar Astrophysics: A Tribute to Kam-Ching Leung*, ed. S. J. Murphy & M. S. Bessell (San Francisco: ASP), p. 57
- Marsh, T., Knigge, C., Pretorius, R., Miller-Jones, J., Koerding, E., Sivakoff, G., Woudt, P., & Warner, B. 2013, *Astron. Telegram*, 4704
- Mason, E., & Howell, S. 2003, *A&A*, 403, 699
- Matsui, R., et al. 2009, *PASJ*, 61, 1081
- Matsumoto, K., Nogami, D., Kato, T., & Baba, H. 1998, *PASJ*, 50, 405
- Mattei, J., Ducoty, R., Stanton, R., & Scovill, C. 1985, *IAU Circ.*, 4026
- Matthews, O. M., Speith, R., Wynn, G. A., & West, R. G. 2007, *MNRAS*, 375, 105
- Mayall, M. W. 1946, *Bull. Harv. Coll. Obs.*, 918, 3
- Maza, J., & Gonzalez, L. E. 1983, *IAU Circ.*, 3856
- McAllister, M. J., et al. 2015, *MNRAS*, 451, 4633
- Mendelson, H., Leibowitz, E. M., Brosch, N., & Almozno, E. 1992, *IAU Circ.*, 5509
- Mennickent, R. E., & Tappert, C. 2001, *A&A*, 372, 563
- Mennickent, R. E., Tappert, C., Gallardo, R., Duerbeck, H. W., & Augusteijn, T. 2002, *A&A*, 395, 557
- Meyer, F., & Meyer-Hofmeister, E. 1981, *A&A*, 104, L10
- Meyer, F., & Meyer-Hofmeister, E. 1994, *A&A*, 288, 175
- Meyer, F., & Meyer-Hofmeister, E. 2015, *PASJ*, 67, 52
- Meyer-Hofmeister, E., Meyer, F., & Liu, B. F. 1998, *A&A*, 339, 507
- Mitrofanova, A. A., Borisov, N. V., & Shimansky, V. V. 2014, *Astrophys. Bull.*, 69, 82
- Mroz, P., et al. 2013, *Acta Astron.*, 63, 135
- Mukadam, A. S., et al. 2010, *ApJ*, 714, 1702
- Mukai, K., et al. 1990, *MNRAS*, 245, 385
- Nakagawa, S., Noguchi, R., Iino, E., Ogura, K., Matsumoto, K., Arai, A., Isogai, M., & Uemura, M. 2013, *PASJ*, 65, 70
- Nakano, S. 2010, *Cent. Bur. Electron. Telegrams*, 2275, 2
- Nakano, S., et al. 2004, *IAU Circ.*, 8363
- Nakano, S., Nishimura, H., Kadota, K., & Yusa, T. 2011a, *Cent. Bur. Electron. Telegrams*, 2616
- Nakano, S., Nishimura, H., Noguchi, T., & Munari, U. 2011b, *Cent. Bur. Electron. Telegrams*, 2818, 1
- Nakano, S., & Noguchi, T. 2012, *Cent. Bur. Electron. Telegrams*, 3229
- Nakata, C., et al. 2014, *PASJ*, 66, 116
- Nakata, C., Kuin, P., Kato, T., & Ohshima, T. 2013a, *Astron. Telegram*, 5253
- Nakata, C., et al. 2013b, *PASJ*, 65, 117
- Naylor, T., Charles, P. A., Hassall, B. J. M., Bath, G. T., Berriman, G., Warner, B., Bailey, J., & Reinsch, K. 1987, *MNRAS*, 229, 183
- Nelemans, G. 2005, in *ASP Conf. Ser. 330, The Astrophysics of Cataclysmic Variables and Related Objects*, ed. J.-M. Hameury, & J.-P. Lasota (San Francisco: ASP), p. 27
- Nesci, R., Caravano, A., Falasca, V., & Villani, L. 2013, *IBVS*, 6059
- Neustroev, V., & Sjöberg, G. 2013, *Astron. Telegram*, 4744
- Nogami, D., Baba, H., Kato, T., & Novák, R. 1998, *PASJ*, 50, 297
- Nogami, D., et al. 2009, in *ASP Conf. Ser. 404, The Eighth Pacific Rim Conference on Stellar Astrophysics: A Tribute to Kam-Ching Leung*, ed. S. J. Murphy & M. S. Bessell (San Francisco: ASP), p. 52
- Nogami, D., & Iijima, T. 2004, *PASJ*, 56, S163
- Nogami, D., Kato, T., Baba, H., Matsumoto, K., Arimoto, J., Tanabe, K., & Ishikawa, K. 1997, *ApJ*, 490, 840
- Nogami, D., Monard, B., Retter, A., Liu, A., Uemura, M., Ishioka, R., Imada, A., & Kato, T. 2004, *PASJ*, 56, L39
- Novák, R., Vanmunster, T., Jensen, L. T., & Nogami, D. 2001, *IBVS*, 5108
- O'Donoghue, D., Chen, A., Marang, F., Mittaz, J. P. D., Winkler, H., & Warner, B. 1991, *MNRAS*, 250, 363
- Ogilvie, G. I., & Dubus, G. 2001, *MNRAS*, 320, 485
- Ortolani, S., Rafanelli, P., Rosino, L., & Vittone, A. 1980, *A&A*, 87, 31
- Osaki, Y. 1974, *PASJ*, 26, 429
- Osaki, Y. 1985, *A&A*, 144, 369
- Osaki, Y. 1989, *PASJ*, 41, 1005
- Osaki, Y. 1995a, *PASJ*, 47, 47
- Osaki, Y. 1995b, *PASJ*, 47, L25
- Osaki, Y. 1996, *PASP*, 108, 39
- Osaki, Y. 1998, in *ASP Conf. Ser. 137, Wild Stars in the Old West*, ed. S. Howell, E. Kuulkers, & C. Woodward (San Francisco: ASP), p. 334
- Osaki, Y. 2003, *PASJ*, 55, 841
- Osaki, Y., & Kato, T. 2013a, *PASJ*, 65, 50
- Osaki, Y., & Kato, T. 2013b, *PASJ*, 65, 95
- Osaki, Y., & Kato, T. 2014, *PASJ*, 66, 15
- Osaki, Y., & Meyer, F. 2002, *A&A*, 383, 574
- Osaki, Y., & Meyer, F. 2003, *A&A*, 401, 325
- Osaki, Y., Meyer, F., & Meyer-Hofmeister, E. 2001, *A&A*, 370, 488
- Osaki, Y., Shimizu, S., & Tsugawa, M. 1997, *PASJ*, 49, L19
- Paczynski, B., & Sienkiewicz, R. 1981, *ApJ*, 248, L27
- Paczynski, B., & Sienkiewicz, R. 1983, *ApJ*, 268, 825
- Patterson, J. 1980, *ApJ*, 241, 235
- Patterson, J. 1998, *PASP*, 110, 1132
- Patterson, J. 2011, *MNRAS*, 411, 2695
- Patterson, J., Augusteijn, T., Harvey, D. A., Skillman, D. R., Abbott, T. M. C., & Thorstensen, J. 1996, *PASP*, 108, 748

- Patterson, J., Bond, H. E., Grauer, A. D., Shafter, A. W., & Mattei, J. A. 1993, *PASP*, 105, 69
- Patterson, J., Jablonski, F., Koen, C., O'Donoghue, D., & Skillman, D. R. 1995, *PASP*, 107, 1183
- Patterson, J., et al. 1998, *PASP*, 110, 1290
- Patterson, J., McGraw, J. T., Coleman, L., & Africano, J. L. 1981, *ApJ*, 248, 1067
- Patterson, J., et al. 2002, *PASP*, 114, 721
- Patterson, J., Richman, H., Kemp, J., & Mukai, K. 1998, *PASP*, 110, 403
- Patterson, J., Schwartz, D. A., Pye, J. P., Blair, W. P., Williams, G. A., & Caillault, J.-P. 1992, *ApJ*, 392, 233
- Patterson, J., et al. 2003, *PASP*, 115, 1308
- Pavlenko, E. 2009, *J. of Physics Conference Series*, 172, 012071
- Pavlenko, E., Antoniuk, O., Antoniuk, K., Andreev, M., Samsonov, D., Baklanov, A., & Golovin, A. 2010, in *AIP Conference Proceedings*, 17th European White Dwarf Workshop, ed. K. Werner & T. Rauch (Maryland: AIP), p. 332
- Pavlenko, E., Kato, T., Sosnovskij, A. A., Andreev, M. V., Ohshima, T., Sklyanov, A. S., Bikmaev, I. F., & Galeev, A. I. 2014, *PASJ*, 66, 113
- Pavlenko, E., et al. 2012, *Mem. Soc. Astron. Ital.*, 83, 520
- Pavlenko, E., et al. 2007, in *ASP Conf. Ser.* 372, 15th European Workshop on White Dwarfs, ed. R. Napiwotzki, & M. R. Burleigh (San Francisco: ASP), p. 511
- Pavlov, M. V., & Shugarov, S. Yu. 1985, *Astron. Tsirk.*, 1373, 8
- Pesch, P., & Sanduleak, N. 1987, *IBVS*, 2989
- Pojmański, G. 2002, *Acta Astron.*, 52, 397
- Price, A., et al. 2004, *PASP*, 116, 1117
- Prieto, J. L., et al. 2013, *Astron. Telegram*, 4999
- Prieto, J. L., et al. 2014a, *Astron. Telegram*, 6475, 1
- Prieto, J. L., et al. 2014b, *Astron. Telegram*, 6340
- Pych, W., & Olech, A. 1995, *Acta Astron.*, 45, 385
- Quimby, R., et al. 2005, *Astron. Telegram*, 658
- Rappaport, S., Joss, P. C., & Webbink, R. F. 1982, *ApJ*, 254, 616
- Richter, G. A. 1968, *IBVS*, 293
- Richter, G. A. 1982, *Comm. of the Konkoly Obs.*, 83, 201
- Richter, G. A. 1985, *IBVS*, 2714
- Richter, G. A. 1986a, *Astron. Nachr.*, 307, 221
- Richter, G. A. 1986b, *IBVS*, 2971
- Richter, G. A. 1991, *IBVS*, 3619
- Richter, G. A. 1992, in *Reviews in Modern Astronomy 5: Variabilities in Stars and Galaxies*, ed. G. Klare (Berlin: Springer), p. 26
- Richter, G. A. 1992, in *ASP Conf. Ser.* 29, *Viña del Mar Workshop on Cataclysmic Variable Stars*, ed. N. Vogt (San Francisco: ASP), p. 12
- Robertson, J. W., Honeycutt, R. K., Hillwig, T., Jurcevic, J. S., & Henden, A. A. 2000, *AJ*, 119, 1365
- Robertson, J. W., Honeycutt, R. K., & Turner, G. W. 1995, *PASP*, 107, 443
- Rogoziecki, P., & Schwarzenberg-Czerny, A. 2001, *MNRAS*, 323, 850
- Romano, G. 1964, *Mem. Soc. Astron. Ital.*, 35, 101
- Samus, N. N. 1982a, *Astron. Tsirk.*, 1205, 2
- Samus, N. N. 1982b, *Astron. Tsirk.*, 1215, 8
- Samus, N. N., V., Durlevich O., Kazarovets, E. V., Kireeva, N. N., Pastukhova, E. N., Zharova, A. V., & et al. 2011, *CDS B/gcvs*
- Savoury, C. D. J., et al. 2011, *MNRAS*, 415, 2025
- Schaefer, B. E., & Hoffleit, D. 1994, *IBVS*, 4123
- Schmeer, P., Hurst, G. M., Kilmartin, P. M., & Gilmore, A. C. 1992, *IAU Circ.*, 5502
- Schmidtobreick, L., Galli, L., Whiting, A., & Tappert, C. 2004, *IBVS*, 5508
- Schoembs, R., & Vogt, N. 1980, *A&A*, 91, 25
- Schwieterman, E. W., et al. 2010, *J. of the Southeastern Assoc. for Research in Astron.*, 3, 6
- Senziani, F., Skinner, G., & Jean, P. 2008, *Astron. Telegram*, 1372
- Shafter, A. W. 2013, *AJ*, 145, 117
- Shafter, A. W., et al. 2009, *American Astron. Soc. Meeting Abstracts*, 213, 491.08
- Shappee, B. J., et al. 2014, *ApJ*, 788, 48
- Sharov, A. S., & Alksnis, A. K. 1989, *Soviet Astronomy Letters*, 15, 382
- Shears, J., et al. 2010, *J. Br. Astron. Assoc.*, 120, 43
- Shears, J., Boyd, D., Krajci, T., Koff, R., Thorstensen, J. R., & Poyner, G. 2008, *J. Br. Astron. Assoc.*, 118, 95
- Shears, J., et al. 2009, *J. Br. Astron. Assoc.*, 119, 340
- Shears, J., Klingenberg, G., & de Ponthiere, P. 2007, *J. Br. Astron. Assoc.*, 117, 331
- Shugarov, S. Y., Volkov, I. M., & Chochol, D. 2008, *IBVS*, 5862
- Shurpakov, S., et al. 2013, *Astron. Telegram*, 5196
- Silvestri, N. M., Szkody, P., Mukadam, A. S., Hermes, J. J., Seibert, M., Schwartz, R. D., & Harpe, E. J. 2012, *AJ*, 144, 84
- Skidmore, W., Mason, E., Howell, S. B., Ciardi, D. R., Littlefair, S., & Dhillon, V. S. 2000, *MNRAS*, 318, 429
- Skidmore, W., Welsh, W. F., Wood, J. H., Catalán, M. S., & Horne, K. 1999, *MNRAS*, 310, 750
- Smak, J. 1993, *Acta Astron.*, 43, 101
- Smak, J. 2004, *Acta Astron.*, 54, 221
- Smak, J. 2013, *Acta Astron.*, 63, 109
- Smak, J. I. 1991, *Acta Astron.*, 41, 269
- Smak, J. I. 2000, *Acta Astron.*, 50, 399
- Smith, D. A., et al. 2002, *IBVS*, 5226
- Soejima, Y., Imada, A., Nogami, D., Kato, T., & Monard, L. A. G. 2009, *PASJ*, 61, 395
- Solheim, J.-E. 2010, *PASP*, 122, 1133
- Southworth, J., et al. 2008, *MNRAS*, 391, 591
- Stanek, K. Z., et al. 2014a, *Astron. Telegram*, 6233
- Stanek, K. Z., et al. 2014b, *Astron. Telegram*, 6061
- Steehghs, D., Harlaftis, E. T., & Horne, K. 1997, *MNRAS*, 290, 28P
- Steehghs, D., Marsh, T., Knigge, C., Maxted, P. F. L., Kuulkers, E., & Skidmore, W. 2001, *ApJ*, 562, L145
- Still, M., Howell, S. B., Wood, M. A., Cannizzo, J. K., & Smale, A. P. 2010, *ApJ*, 717, L113
- Szkody, P. 1987, *ApJS*, 63, 685
- Szkody, P., Desai, V., & Hoard, D. W. 2000, *AJ*, 119, 365
- Szkody, P., et al. 2003, *AJ*, 126, 1499
- Szkody, P., et al. 2006, *AJ*, 131, 973
- Szkody, P., Howell, S. B., Mateo, M., & Kreidl, T. J. 1989, *PASP*, 101, 899
- Szkody, P., et al. 2010, *ApJ*, 710, 64
- Szkody, P., et al. 2007, *ApJ*, 658, 1188
- Szkody, P., Mukadam, A. S., Sion, E. M., Gansicke, B. T., Henden, A., & Townsley, D. 2013, *AJ*, 145, 121
- Templeton, M., et al. 2007, *Cent. Bur. Electron. Telegrams*, 1053
- Templeton, M. R. 2009, *AAVSO Alert Notice*, 407
- Templeton, M. R., et al. 2006, *PASP*, 118, 236
- Teyssier, F. 2014, *Astron. Telegram*, 6235

- Thiele, H. 1916, *Astron. Nachr.*, 202, 213
- Thorstensen, J. R. 2003, *AJ*, 126, 3017
- Thorstensen, J. R., Patterson, J. O., Kemp, J., & Vennes, S. 2002, *PASP*, 114, 1108
- Thorstensen, J. R., & Skinner, J. N. 2012, *AJ*, 144, 81
- Tovmassian, G., Clark, D., & Zharikov, S. 2010, *Cent. Bur. Electron. Telegrams*, 2283, 1
- Uemura, M., et al. 2008a, *PASJ*, 60, 227
- Uemura, M., et al. 2008b, *IBVS*, 5815, 1
- Uemura, M., Kato, T., Ohshima, T., & Maehara, H. 2012, *PASJ*, 64, 92
- Uemura, M., Nogami, D., & Kato, T. 2015, *PASJ*, submitted
- Uthas, H., et al. 2012, *MNRAS*, 420, 379
- van Amerongen, S., Bovenschen, H., & van Paradijs, J. 1987, *MNRAS*, 229, 245
- van der Woerd, H., van der Klis, M., van Paradijs, J., Beuermann, K., & Motch, C. 1988, *ApJ*, 330, 911
- van der Woerd, H., & van Paradijs, J. 1987, *MNRAS*, 224, 271
- van Teeseling, A., Hessman, F. V., & Romani, R. W. 1999, *A&A*, 342, L45
- van Zyl, L., et al. 2004, *MNRAS*, 350, 307
- van Zyl, L., Warner, B., O'Donoghue, D., Sullivan, D., Pritchard, J., & Kemp, J. 2000, *Baltic Astron.*, 9, 231
- Vanmunster, T., et al. 2006, *Society for Astronom. Sciences Ann. Symp.*, 25, 77
- Vanmunster, T., & Sarneczky, K. 1997, *IAU Circ.*, 6740
- Vican, L., et al. 2011, *PASP*, 123, 1156
- Vladimirov, V., et al. 2014, *Astron. Telegram*, 6059
- Vladimirov, V., et al. 2013, *Astron. Telegram*, 5481
- Vogt, N. 1983, *A&A*, 118, 95
- Vogt, N., & Bateson, F. M. 1982, *A&AS*, 48, 383
- Waagen, E. O., Schmeer, P., Stubbings, R., & Pearce, A. 2007, *IAU Circ.*, 8829
- Warner, B. 1987, *MNRAS*, 227, 23
- Warner, B. 1995, *Cataclysmic Variable Stars* (Cambridge: Cambridge University Press)
- Warner, B., Livio, M., & Tout, C. A. 1996, *MNRAS*, 282, 735
- Warner, B., & Pretorius, M. L. 2008, *MNRAS*, 383, 1469
- Warner, B., & van Zyl, L. 1998, in *IAU Symp.* 185, *New Eyes to See Inside the Sun and Stars*, ed. F.-L. Deubner, J. Christensen-Dalsgaard, & D. Kurtz (Berlin: Springer), p. 321
- Warner, B., & Woudt, P. A. 2004, in *ASP Conf. Ser.* 310, *IAU Colloq. 193: Variable Stars in the Local Group*, ed. D. W. Kurtz & K. R. Pollard (San Francisco: ASP), p. 382
- Weber, R. 1966, *IBVS*, 123
- Wenzel, W. 1983, *Mitteil. Veränderl. Sterne*, 9, 141
- Wenzel, W. 1991a, *IBVS*, 3626
- Wenzel, W. 1991b, *IBVS*, 3689
- Wild, P. 1979, *IAU Circ.*, 3412
- Williams, S. C., Darnley, M. J., Bode, M. F., & Copperwheat, C. M. 2015, *Astron. Telegram*, 6992
- Wils, P., Gänsicke, B. T., Drake, A. J., & Southworth, J. 2010, *MNRAS*, 402, 436
- Wolf, M. 1919, *Astron. Nachr.*, 209, 85
- Wolf, M., & Wolf, G. 1905, *Astron. Nachr.*, 169, 415
- Wood, M. A., Still, M. D., Howell, S. B., Cannizzo, J. K., & Smale, A. P. 2011, *ApJ*, 741, 105
- Woudt, P. A., & Warner, B. 2001, *MNRAS*, 328, 159
- Woudt, P. A., & Warner, B. 2002a, *Ap&SS*, 282, 433
- Woudt, P. A., & Warner, B. 2002b, *MNRAS*, 335, 44
- Woudt, P. A., & Warner, B. 2011, *Ap&SS*, 333, 119
- Woudt, P. A., Warner, B., & Pretorius, M. L. 2004, *MNRAS*, 351, 1015
- Wu, X., Li, Z., Gao, W., & Leung, K. 2001, *ApJ*, 549, L81
- Yamaoka, H. 2010, *Cent. Bur. Electron. Telegrams*, 2273
- Yamaoka, H., et al. 2011, *Cent. Bur. Electron. Telegrams*, 2731, 1
- Yamaoka, H., Itagaki, K., Korotkiy, S., & Samus, N. N. 2008a, *IAU Circ.*, 8971
- Yamaoka, H., Itagaki, K., Martignoni, M., & Koff, R. A. 2012, *Cent. Bur. Electron. Telegrams*, 3228
- Yamaoka, H., Itagaki, K., Naito, H., & Narusawa, S. 2008b, *Cent. Bur. Electron. Telegrams*, 1216
- Zemko, P. O., & Kato, T. 2013, *Astrophysics*, 56, 203
- Zharikov, S., Tovmassian, G., Aviles, A., Michel, R., Gonzalez-Buitrago, D., & Garcia-Diaz, M. T. 2013, *A&A*, 549, A77
- Zharikov, S. V., Tovmassian, G. H., Napiwotzki, R., Michel, R., & Neustroev, V. 2006, *A&A*, 449, 645
- Zharikov, S. V., et al. 2008, *A&A*, 486, 505
- Zheng, W., et al. 2010, *Cent. Bur. Electron. Telegrams*, 2574